

Utah State University

DigitalCommons@USU

Aspen Bibliography

Aspen Research

1979

Equations Predicting Primary Productivity (Biomass) of Trees, Shrubs and Lesser Vegetation Based on Current Literature

W. Stanek

D. State

Follow this and additional works at: https://digitalcommons.usu.edu/aspen_bib



Part of the [Agriculture Commons](#), [Ecology and Evolutionary Biology Commons](#), [Forest Sciences Commons](#), [Genetics and Genomics Commons](#), and the [Plant Sciences Commons](#)

Recommended Citation

Stanek, W. and State, D. 1979. Equations predicting primary productivity (biomass) of trees, shrubs and lesser vegetation based on current literature. Environment Canada, Forestry Service. Pacific Forest Research Centre, BC-X, 183, Victoria B.C.

This Report is brought to you for free and open access by the Aspen Research at DigitalCommons@USU. It has been accepted for inclusion in Aspen Bibliography by an authorized administrator of DigitalCommons@USU. For more information, please contact digitalcommons@usu.edu.



LOAN COPY

4-79

Equations Predicting
Primary Productivity
(Biomass) of Trees,
Shrubs and Lesser
Vegetation Based on
Current Literature

W. Stanek
D. State

PSW/ FOREST AND RANGE
EXPERIMENT STATION

JAN 4 1979

STATION LIBRARY COPY



Environment
Canada

Environnement
Canada

Forestry
Service

Service
des Forêts

ABSTRACT

The report contains regression equations for calculations of biomass of several mainly Canadian forest species (trees, shrubs, lesser vegetation) and their components (leaves, branches, bole, bark, wood, roots, etc.).

The lists are arranged by plant species in alphabetical order of common names. Glossaries and a sketch define the tree components. Many pertinent references are given.

RÉSUMÉ

Ce rapport contient des équations de régression pour le calcul de la biomasse de plusieurs espèces forestières surtout canadiennes (arbres, arbustes, végétation basse) et leurs parties (feuilles, branches, fût, écorce, bois, racines, etc.).

Sont énumérés les noms communs des plantes par ordre alphabétique. Des glossaires et un dessin définissent les parties des arbres. Maintes références pertinentes sont incluses.

INTRODUCTION

Forest productivity may be envisaged from two distinct aspects: economic and biological (Duvigneaud in Anon. 1971b). The former evaluates the volume of solid wood per unit area, the latter the dry weight of tree components in kilocalories. Economic forest productivity is familiar to most foresters and has long been an essential part of forest management. In contrast, biological forest productivity has come into focus relatively recently, as a result of advances in knowledge of the ecosystem (biogeocoenosis) or concern with energy supply.

The foundations of the science of biological productivity of forests were laid by Boysen Jensen (1910 and 1932), who defined the concepts and developed techniques and methods, though, in the last century, considerable contributions were made by pioneers such as Ebermayer in Germany or Henry in France. Among other works, Ovington's (1957) "Dry matter production of *Pinus sylvestris*" is a classic, and Lieth's (1962) Proceedings of the Symposium "Stoffproduktion der Pflanzendecke" (productivity of the vegetal cover), held in Stuttgart-Hohenheim, is important in that it identifies problems and methods.

The impetus for more research came in 1963 when, under the auspices of the International Biological Program (IBP), the "Section PT" (Productivity of Terrestrial Communities) met and proposed a definitive 7-year program (Duvigneaud in Anon. 1971b).

Since then, several IBP meetings have taken place and many findings have been published. Those accessible to us have been listed under References. Of special interest are the Proceedings of the Brussels Symposium in October 1969 (Anon. 1971b), which is actually a 6-year IBP progress report and state-of-art review of productivity of forest ecosystems.

In the latter report, primary biological forest productivity (PBFP)^{1/} is defined by Duvigneaud as the rate of production by which energy is stored through the activity of producing organisms, photosynthetic or chemosynthetic, in the form of an organic substance suitable for alimentary utilization.

The gross PBFP (total photosynthesis, total assimilation) is the rate of total photosynthesis, including the organic matter utilized for respiration during the period of measuring. The net PBFP (apparent photosynthesis, net assimilation) is the rate of organic matter production in the form of vegetable tissues, less the matter utilized for respiration during the period of measuring. The followers of Boysen Jensen define the net PBFP as the gross dry matter production minus the dry matter of lost leaves, branches, roots, and through respiration of leaves, branches, stem and roots. It is important to note the basic difference between the two definitions.

The net primary biological productivity may also be defined as the rate of biomass production (energy storage). Newbould (1967) defined biomass as the total amount of photosynthetic plants making up a woodland stand and takes it to include heartwood and bark (which may no longer be alive) but not dead roots and dead branches. In context, biomass should be expressed in terms of dry weight, or ash-free dry weight (=organic weight), and can be estimated directly by weighing or indirectly from measurements of the volume and density of the various components concerned.

The fundamentals for estimation of biomass and its increment, as well as field sampling methods, treatment of samples and evaluation of data have been presented in many publications. Among them are those by Boysen Jensen 1932; Kira et al. 1967; Newbould 1967; Milner and Hughes 1968; Duvigneaud in Anon. 1971b, pp. 111-140; Whittaker and Woodwell in Anon. 1971b, pp. 159-175.

Of prime interest is the determination of periodic biomass increment ΔB during the time period Δt or annual biomass increment (when the period amounts to 1 year).

$$\Delta B = B_2 - B_1 = P_n - (L + G)$$

Where B_2 and B_1 represent the biomass at the end and beginning of the period, P_n stands for net productivity during Δt , L for plant losses by death and shedding during Δt and G for plant losses owing to consumer during Δt .

Biomass data are essential for the quantita-

^{1/} The secondary productivity is the rate of energy accumulation at trophic levels of consumers and decomposers, and obviously will depend on vegetable matter produced by primary productivity.

tive evaluation of renewable resource potential (Keays in Anon. 1971). They have been used for comparing plant communities, studying the biological and physical processes that affect productivity, and energy flow and utilization relationships in nature. They are also important in forest fuel estimations, and in determining fire behavior in terms of both wildfire and prescribed burning on forest lands (Brown 1965; Kiil 1967; Kurucz 1969; and others).

Biomass determinations may also provide estimates of pulp yield and a basis for determining the feasibility of complete tree utilization for such purposes as wood products, ruminant fodder (Reid 1976; Young 1976; and others), chemicals and fuel.

Fresh or oven-dry biomass of trees, shrubs or lesser vegetation and the components thereof, such as foliage, branches, roots and tree bole, etc., can be determined by many destructive and non-destructive methods. In studies referred to, relationships were developed of biomass and easily determinable characteristics (diameter at breast height or at ground level, tree height, etc.) to facilitate practical application.

The objective of this report is to compile, for use at the PFRC, from available literature, regression equations for calculation of biomass of forest communities and individual plants (trees, shrubs, lesser vegetation) and their components (leaves, branches by size class, bole, merchantable bole, bark, wood, roots by size classes, etc.).

METHODS

Most of the publications concerned with biomass, contain information on methods used by the authors. It is not intended to review the many approaches and techniques; instead, a list is presented of regression equations found in the literature (Appendix). The occurring variables, symbols of which have been standardized, are defined in the "Glossary of Independent Variables". The tree components (Fig. 1) for which the biomass equations were developed, have also been given in standardized form and are defined in the "Glossary of Equation Titles". The glossaries are self explanatory and refer to notations used in the Appendix.

The lists of regression equations (Appendix) are arranged by plant species in alphabetical order of

common names. In the first column are given the species, author, year of publication and equation title. Question mark (?) in the equation title indicates that this information was not given in the original report. Biomass fresh weights (FW) or oven-dry weights (ODW) are, with two exceptions, per individual trees, and are shown in column "FW or ODW". The published regression equations are given in the column "Prediction Equations". The numbers of observations on which the equations are based are contained in column "n" and the coefficient of determination (R^2) or the simple correlation coefficient (r) under column " R^2 or r ". Where (r) is not specified, values are taken as denoting R^2 . In all cases, R^2 and r values were rounded up to two decimals except where they approached 1.0, in which instances they were taken as .99. The range of diameters at ground level, or diameters at breast height, is shown in the last column as "Range of diameters at ground level in mm or diameters at breast height in cm".

Scientific names of plant species are given in the "Glossary of Species Names".

Glossary of Equation Titles

The names of variables used by the different authors have been standardized and defined below to facilitate comparisons. The individual tree components are shown in Figure 1 (based on Young et al. 1964), and are defined as follows.

- 1) Small roots - all roots or root portions as a general rule less than 2.5 cm (1") in diameter.
- 2) Medium roots - roots as a general rule from 2.5-10 cm (1-4") in diameter.
- 3) Large roots - roots as a general rule larger than 10 cm (4").
- 4a) Stump above ground level.
- 4b) Stump below ground - usually to where large roots are apparent.
- 5) Bole - merchantable stem from the stump to a merchantable top diameter.
- 6) Large branches - usually larger than 2.5 cm (1").

7a) Small branches - usually smaller than 2.5 cm (1").

7b) Leaves or needles.

8) Unmerchantable top.

The equations for individual components or groups of components have titles that can be defined in the above terms as follows:

The following additional abbreviations were used mainly to save space: > = larger than, < = smaller than, + = plus, - = minus, & = and, " = inches, cm = centimeters, mm = millimeters, lbs = weight in pounds, kg = weight in kilograms, g = weight in grams, Dom = dominant, Codo = codominant, t = top diameter of stem, s = stump height.

In the following cases, the dependent variables were not precisely defined: Ribe (1973) did not

Equation Title	Dependent variables	Components (shown in Fig. 1)
Tree tot.	Total or complete tree or plant	1,2,3,4,5,6,7,8
Tree	Tree or whole plant above ground level	4a,5,6,7,8
Tree a.s.* * known stump height	Tree above stump	5,6,7,8
Stem	Stem above ground level	4a,5,8
Stem a.s.* * known stump height	Stem above stump	5,8
Bole s* t** * known stump height ** known top diameter	Merchantable stem	5
Stump	Stump	4a+b
Roots	Roots	1,2,3
Crown	Branches & foliage	6,7a+b
Branch	Branches	6,7a
Foliage	Needles or leaves	7b
Bark	Bark only	
Wood	Wood only	
Slash t** ** known top diameter	Branches, foliage & top	6, 7a+b,8

All components and diameter measurements are understood to include bark (outside bark, o.b.) unless stated as inside bark (i.b.). Pertinent sizes of branches and roots are indicated in the equation title.

define stem, but probably means stem above ground level. Kurucz (1969) did not define bole, but is probably equivalent to stem.




Figure 1. Diagrammatic sketch showing the components of a tree.
(adapted from Young et al. 1964)

GLOSSARY OF INDEPENDENT VARIABLES

List of notations used in Appendix. All diameters were measured outside bark.

A_1	= Total age in years	D_8	= (D^2H) = diameter at breast height squared (cm ²) height (cm)
A_2	= $1/A_1$	D_9	= (D^2H) = diameter at breast height squared (cm ²) height (m)
A_3	= Reciprocal of increment core specific gravity	DC_1	= $(D \times CLE)$ Diameter at breast height (in) multiplied by crown length (ft)
BA	= Basal area at breast height $r^2 \pi$ (cm ²)	DC_2	= $(D \times CLE)^2$ The product to the square of diameter at breast height (in) and crown length (ft)
BR	= Branchiness class (1-5)	DC_3	= $(D \times CW^2)$ diameter at breast height (in) multiplied by crown width squared (ft ²)
CH	= Canopy height (m)	g/m^2	= Grams per meter squared
CL	= Crown length (cm)	ha	= Hectare
CL_m	= Crown length (m)	H_1	= Height above ground (ft)
CR	= Crown ratio (CL/Height x 100%)	H_2	= Height above ground (cm)
CS	= Crown surface = $\pi/4 CW \sqrt{CL^2 + CW^2}$	H_3	= Height above ground (m)
CV	= Crown volume $\pi/12(CW^2)CL$	IV	= $(D_2^2 H_3)/100$
CW	= Crown width (ft)	L	= Geographic latitude
CW_m	= Crown width (m)	\log	= Common logarithm
D	= Diameter at crown base (cm) - just below leaf bearing branch	\ln	= Natural logarithm
D_1	= Diameter at breast height (in)	ME	= Average diffuse light intensity (micro-einstein m ² sec ⁻¹ = 6.02×10^{17} photons)
D_2	= Diameter at breast height (cm)	MT	= Metric tons
D_3	= Diameter at breast height (m)	n	= Number of observations
D_4	= Diameter at ground level (mm)	P	= Per cent
D_5	= Diameter at ground level (cm)	r	= Simple correlation coefficient
D_{11}	= Diameter at ground level (in)	R^2	= Coefficient of determination
D_6	= (D^2H) = diameter at breast height squared (in ²) height (ft)	S_1	= Stump diameter (cm)
D_7	= (D^2H) = diameter at breast height squared (m ²) height (m)	SI	= Site Index at age 50
		TA	= Trees per acre
		V	= % vegetation cover

GLOSSARY OF SPECIES NAMES

Alder - Speckled	- <u>Alnus rugosa</u> (DuRoi) Spreng.
- Red	- <u>Alnus rubra</u> Bong.
Aspen - Trembling (Quaking)	- <u>Populus tremuloides</u> Mich.
Bayberry	- <u>Myrica pensylvanica</u> Loisel
Beech	- <u>Fagus grandifolia</u> Ehrh.
Birch - Gray	- <u>Betula populifolia</u> Marsh.
- White (paper)	- <u>Betula papyrifera</u> Marsh.
- Yellow	- <u>Betula alleghaniensis</u> Britt.
Blueberry - Lowbush	- <u>Vaccinium vacillans</u> Torr.
Cedar - Northern White	- <u>Thuja occidentalis</u> L.
- Western Red	- <u>Thuja plicata</u> Donn
Cherry - Choke	- <u>Prunus virginiana</u> L.
- Pin	- <u>Prunus pensylvanica</u> L.f.
Douglas-fir - see Fir - Douglas	
Fern - Sweet	- <u>Comptoria peregrina</u> (L.) Coult.
- Bracken	- <u>Pteridium aquilinum</u> (L.) Kuhn.
Fir - Amabilis (Pacific Silver)	- <u>Abies amabilis</u> (Dougl.) Forb.
- Balsam	- <u>Abies balsamea</u> (L.) Mill.
- Douglas	- <u>Pseudotsuga menziesii</u> (Mirb.) Franco
- Subalpine	- <u>Abies lasiocarpa</u> (Hook.) Nutt.
Hemlock - Eastern	- <u>Tsuga canadensis</u> (L.) Carr
- Mountain	- <u>Tsuga mertensiana</u> (Bong.) Carr.
- Western	- <u>Tsuga heterophylla</u> (Raf.) Sarg.
Hobblebush	- <u>Viburnum alnifolium</u> Marsh.
Honeysuckle - American Fly	- <u>Lonicera canadensis</u> Bartr.
Holly - Canada	- <u>Ilex verticillata</u> (L.) A. Gray
- False	- <u>Nemopanthus mucronata</u> Trel.
Huckleberry - Black	- <u>Gaylussacia baccata</u> Wang.
Inkberry	- <u>Ilex glabra</u>
Labrador Tea	- <u>Ledum groenlandicum</u> Oedr.
Laurel - Sheep	- <u>Kalmia augustifolia</u> (L.)
Leatherleaf	- <u>Chamaedaphne calyculata</u> (L.) Moench.

Maple - Red	- <u>Acer rubrum</u> (L.)
- Striped	- <u>Acer pensylvanicum</u> (L.)
- Sugar	- <u>Acer saccharum</u> March.
Oak - Black	- <u>Quercus velutina</u> Lam.
- Red	- <u>Quercus rubra</u> (L.)
- Scarlet	- <u>Quercus coccinea</u> Muenchh.
- Scrub	- <u>Quercus ilicifolia</u> Wang.
- White	- <u>Quercus alba</u> (L.)
Pine- Jack	- <u>Pinus banksiana</u> Lamb.
- Lödgpole	- <u>Pinus contorta</u> Dougl.
- Pitch	- <u>Pinus rigida</u> Mill.
- Red	- <u>Pinus resinosa</u> Ait.
- Scots	- <u>Pinus sylvestris</u> (L.)
- Slash	- <u>Pinus elliotii</u> Engelm.
- White	- <u>Pinus strobus</u> L.
Rhododendron	- <u>Rhododendron canadense</u> Marsh.
Salal	- <u>Gaultheria shallon</u> Pursh.
Spiraea - meadowsweet	- <u>Spiraea</u> spp. (Tourn.) L.
Spruce - Black	- <u>Picea mariana</u> (Mill.) B.S.P.
- Norway	- <u>Picea abies</u> (L.) Karst.
- Red	- <u>Picea rubens</u> Sarg.
- White	- <u>Picea glauca</u> (Moench) Voss
Willow	- <u>Salix</u> spp. (Tourn.) L.
Witch Hazel	- <u>Hamamelis virginiana</u> (L.)
Whithe - Rod	- <u>Viburnum cassinoides</u> (L.)

Species names for trees are according to: Harlow & Harrar 1958.

Species names for lesser vegetation are according to: Peck 1941 & Marie-Victorin 1964.

REFERENCES

Under references are included all literature perused in search of publications containing information on prediction equations, as well as that specifically mentioned in the text and Appendix.

- Adams, E.L. 1971. Effect of moisture loss on red oak sawlog weight. USDA, For. Serv., Northeast For. Exp. Sta., Upper Darby, Pa., Res. Note NE-133: 4pp.
- Anonymous. 1966. Native trees of Canada. Can. Dept. of Forest and Rural Development, Bull. 61: 291pp.
- Anonymous. 1968. Functioning of terrestrial ecosystems at the primary production level. Proceedings of the Copenhagen Symposium. Ed. F.E. Eckardt. Natural Resources Ser. Vol. IV. Unesco, Place de Fontenoy, 75 Paris 7e. 516pp.
- Anonymous. 1970. Productivity and conservation in northern circumpolar lands. Proceedings of a conference Edmonton, Alberta, October 1969. IUCN Publications new series No. 16. Ed. W.A. Fuller and P.G. Kevan, Morges, Switzerland. 344pp.
- Anonymous. 1971a. Forest biomass studies. IUFRO Section 25, Yield and Growth, Working Group on Forest Biomass Studies, Life Sciences and Agriculture, Exp. Sta., Univ. of Maine at Orono, Miscell. Publ. 132, 205pp.
- Anonymous. 1971b. Productivity of forest ecosystems. Proc. Brussels Symp. UNESCO and IBP October 27-3, 1969. Ed. P. DuVigneaud, UNESCO Place de Fontenoy, 75 Paris - 7e, Ecology and Conservation No. 4: 707pp.
- Banks, C.H. 1974. C-U-T-T or the complete utilization of trees and timber - fact or fantasy. Timber Research Unit, Council for Scientific and Industrial Research, P.O. Box 395, Pretoria, South Africa. Getting more out of our timber resources. 5(s.99): 12pp.
- Baskerville, G.L. 1964. Distribution of dry weight in immature balsam fir trees. Can. Dept. For., For. Res. Branch, Fredericton, N.B., Rep. 64-M-6: 6pp.
- Baskerville, G.L. 1965. Dry matter production in immature balsam fir stands. For. Sci. Monograph 9: 42pp.
- Baskerville, G.L. 1965a. Estimation of dry weight of tree components and total standing crop in conifer stands. Ecology 46(6): 867-869.
- Baskerville, G.L. 1966. Dry-matter production in immature balsam fir stands: roots, lesser vegetation, and total stand. For. Sci. 12(1): 49-53.
- Boysen Jensen, P. 1910. Studien ueber das Verhaeltnis der Waldbaeume zum Licht. (Studies of forest tree and light relationships.) Tidsskr. Skovvaes. 22: 1-116.
- Boysen Jensen, P. 1932. Die Stoffproduktion der Pflanzen. (The matter production of plants.) Fischer, Jena, 108pp.
- Brown, J.K. 1963. Crown weights in red pine plantations. U.S.D.A., For. Serv., Lake States For. Exp. Sta., St. Paul, Minn., Research Note LS-19: 4pp.
- Brown, J.K. 1965. Estimating crown fuel weights of red pine and jack pine. U.S.D.A., For. Serv. Lake States For. Exp. Sta., St. Paul, Minn., Res. Pap. LS-20: 12pp.
- Carlisle, A. 1976. The utilization of forest biomass and forest industry wastes for the production and conservation of energy 1976. Can. Dept. Environ., Can. For. Serv., Petawawa For. Exp. Sta., Chalk River, Ontario: 54pp.
- Chase, J., F. Hyland and H.E. Young. 1971. Puckerbrush pulping studies. Life Sci. and Agri. Exp. Sta., Univ. of Maine at Orono, Techn. Bull. 49: 64pp.
- Chase, J., F.H. Hyland and H.E. Young. 1973. The commercial use of puckerbrush pulp. Life. Sci. and Agr. Exp. Sta., Univ. of Maine at Orono, Tech. Bull. 65: 54pp.
- Chase, J. and E. Young. 1976. Corrugating pulp from puckerbrush. Research in the life sciences. Life. Sci. and Agr. Exp. Sta., Univ. of Maine at Orono 23(10): 9pp.
- Chase, J. and E. Young. 1976. The potential of softwood thinnings and standing dead softwoods as a source of wood pulp. Life Sci. and Agri. Exp. Sta., Univ. of Maine at Orono, Tech. Bull. 82: 25pp.
- Curtis, H. 1965. Tree weight equations - their development and use in forest management planning. Proc. Soc. of Amer. Foresters, Detroit, Michigan: 189-191.
- Davis, C.C. 1963. On questions of production and productivity in ecology. Archeological Hydrobiology. 59(2): 145-161.
- Doerner, K., Jr. 1964. Some causes and effects of horizontal density variation in tree stems. For. Sci. 10(1): 24-27.
- Doucet, R., J.V. Berglund and C.E. Farnsworth. 1976. Dry matter production in 40-year Pinus banksiana stands in Quebec. Can. J. For. Res. 6(3): 357-367.
- Drolet, J.C., R.M. Newnham and T.B. Tsay. 1971. Branchiness of jack pine, black spruce and balsam fir in relation to mechanized delimbing. Can. Dept. Environ., Can. For. Serv., Forest Management Instit. Info. Rep. FMR-X-34: 34pp.

- Dutrow, G.F. 1971. Economic implications of silage sycamore. U.S.D.A., Forest Service, South. For. Exp. Stn., New Orleans, La. Res. Paper SO-67: 9 pp.
- Duvigneaud, P. 1971. Concepts sur la productivité primaire des écosystèmes forestiers. p111-140 in Anon. 1971b. Productivity of forest ecosystems.
- Dyer, F.R. 1967. Fresh and dry weight, nutrient elements and pulping characteristics of northern white cedar *Thuja occidentalis*. School of Forestry, Univ. of Maine. Maine Agr. Exp. Sta. Tech. Bull. 27: 40pp.
- Dyer, F.R., A.J. Chase and H.E. Young 1968. Pulp from presently non-commercial woody perennials. Reprint Pulp and Paper Mag. of Can., January 5, 1968 issue: 6pp.
- Erickson, J.R. 1972. The moisture content and specific gravity of the bark and wood of northern pulpwood species. U.S.D.A., For. Serv., North Centr. For. Exp. Sta., St. Paul, Minn., Res. Note NC-141, 3pp.
- Fahnestock, G.R. and W.K. Key. 1971. Weight of brushy forest fire fuels from photographs. For. Sci. 17(1): 119-124.
- Flann, I.B. 1962. Some conversion factors and related information. Can. Dept. Forestry, For. Prod. Res. Branch, Ottawa Lab., Tech. Note 26: 23pp.
- Fleischer, H.O. 1971. The impact of utilization research on the complete use of the forest. Wood Sci. and Tech. 5: 247-254.
- Forbes, E.W. 1968. How to estimate weight of chips produced from pulpwood. U.S. For. Serv. North Centr. For. Exp. Sta., St. Paul, Minnesota: 2pp.
- Fortesque, J.A.C. and G.G. Marten. 1970. Micronutrients: forest ecology and systems analysis. In Ecological studies. Analysis and Synthesis, Ed. D.E. Reichle, Springer Verlag, Heidelberg, New York, Vol 1: 173-198.
- Fortesque, J.A.C. 1971. Biogeochemistry, plant growth and the environment. The Can. Mining and Metallurgical Bull. Ontario (CIM) Bull. Aug. p. 77-82.
- Fortesque, J.A.C. and J. Hughes. 1971. A preliminary study of the use of branch material for the estimation of the trace element status of a forest stand. Dept. Geological Sci. Brock University, St. Catharines, Ontario. Res. Rep. Ser. No. 1, Studies in Forest Biogeochemistry No. 1. 28pp.
- Gary, H.L. 1976. Crown structure and distribution of biomass in a lodgepole pine stand. U.S.D.A., Forest Service, Rocky Mt. For. and Range Exp. Stn., Fort Collins, Colo. Res. Paper RM-165: 20pp.
- Gerry, R.W., H.E. Young, W.P. Apgar and H.C. Dickey. 1977. Muka feeding trials with Maine farm livestock. Research in Life Sciences, Agri. Exp. Sta., Univ. of Maine at Orono. 24(1-2): 6pp.
- Grier, C.C. and R.H. Waring. 1974. Conifer foliage mass related to sapwood area. For. Sci. 20: 205-206.
- Hakkila, P. 1971. Coniferous branches as a raw material source. Communications Inst. For. Fenniae, Helsinki 75(1): 60pp.
- Hakkila, P. 1972. Mechanized harvesting of stumps and roots. Comm. Inst. For. Fenniae, Helsinki 77(1): 71pp.
- Hale, J.D. and K.G. Fensom. 1931. The rate of growth and density of the wood of white spruce. Dept. of the Interior, Canada For. Serv. Circ. 30: 19pp.
- Hale, J.D. and J.B. Prince. 1940. Density and rate of growth in the spruces and balsam fir of eastern Canada. Canada Dept. of Mines and Resources, Dominion For. Serv. Bull. 94: 43pp.
- Hardy, S.S. and G.W. Weiland. 1964. Weight as a basis for the purchase of pulpwood in Maine. Univ. of Maine at Orono. Maine Agr. Exp. Sta. Tech. Ser., Bull. 14: 64pp.
- Harlow, W.M. and E.S. Harrar. 1958. Textbook of Dendrology. 4th Ed. McGraw-Hill, New York, Toronto, London, 561 pp.
- Hegyi, F. 1969. A study of basic density variation in jack pine. Can. Dept. of Fisheries and Forestry, Can. For. Serv., For. Res. Lab., Sault Ste. Marie, Ontario. Internal Rep. 0-17: 56pp.
- Hegyi, F. 1972. Dry matter distribution in jack pine stands in northern Ontario. Repr. from The For. Chron. 48(4): 5pp.
- Honer, T.G. 1970. Bole weight to volume ratio: a constant for open grown balsam fir. Can. Dept. Fish. For., Can. For. Serv., Bimonthly Res. Notes 26: 37pp.
- Honer, T.G. 1971. Weight relationships in open- and forest-grown balsam fir trees. In Anon. 1971a, Forest Biomass studies. pp. 65-78.
- Horn, R.A. and R.J. Auchter. 1972. Kraft pulping of pulpwood chips containing bark. Paper Trade J. Nov. 6: pp. 55-59.
- Howard, O. and F.R. Ward. 1972. Measurement of logging residue - alternative applications of the line intersect method. Pac. N.W. For. Range Exp. Sta., For. Serv., U.S.D.A., Portland, Oregon. Res. Note PNW-183: 8pp.
- Johnstone, W.D. 1970. Component dry weights of 100-year-old lodgepole pine trees. Can. Dept. of Fisheries and Forestry, Can. For. Serv., For. Res. Lab., Calgary, Alberta. Info Rep. A-X-31: 14pp.
- Johnstone, W.D. 1970a. Some variations in specific gravity and moisture content of 100-year-old lodgepole pine trees. Can. Dept. of Fisheries and Forestry, Can. For. Serv., For. Res. Lab., Calgary, Alberta. Info. Rep. A-X-29: 19pp.

- Keays, J.L. 1968. Whole-tree utilization studies; selection of tree components for pulping research. Can. Dept. Fisheries and Forestry, Can. For. Serv., Forest Products Lab., Vancouver, B.C. Info. Rep. VP-X-35: 31pp.
- Keays, J.L. 1971. Complete-tree utilization, an analysis of the literature. Part I: unmerchantable top of bole. Can. Dept. Fisheries and Forestry, Can. For. Serv., Forest Products Lab., Vancouver, B.C. Info. Rep. VP-X-69: 98pp.
- Keays, J.L. 1971. Complete-tree utilization, an analysis of the literature. Part II: Foliage. Can. Dept. Fisheries and Forestry, Can. For. Serv., Forest Products Lab., Vancouver, B.C. Info. Rep. VP-X-70: 94pp.
- Keays, J.L. 1971. Complete-tree utilization, an analysis of the literature. Part III: Branches. Can. Dept. Fisheries and Forestry, Can. For. Serv., Forest Products Lab., Vancouver, B.C. Info. Rep. VP-X-71: 67pp.
- Keays, J.L. 1971. Complete-tree utilization, an analysis of the literature. Part IV: Crown and slash. Can. Dept. Fisheries and Forestry, Can. For. Serv., Forest Products Lab., Vancouver, B.C. Info. Rep. VP-X-77: 79pp.
- Keays, J.L. 1971. Complete-tree utilization, an analysis of the literature. Part V: Stump, Roots and stump-root system. Can. Dept. Fisheries and Forestry, Can. For. Serv., Forest Products Lab., Vancouver, B.C. Info. Rep. VP-X-79: 62pp.
- Keays, J.L. 1971. Complete-tree utilization, Resume of a literature Review. In Anon, 1971, Forest Biomass Studies. pp 93-102.
- Keays, J.L. and K. Hunt. 1971. Departmental pulping service studies. Can. Dept. Fisheries and Forestry, Can. For. Serv., Forest Products Lab., Vancouver, B.C. Internal Report. 60pp.
- Keen, R.E. 1963. Weights and centres of gravity involved in handling pulpwood trees. Pulp and paper Res. Instit. of Canada, Montreal, Canada. Tech. Rep. No. 340 (Woodlands Res. Index No. 147): 93pp.
- Kiil, A.D. 1967. Fuel weight tables for white spruce and lodgepole pine crowns in Alberta. Can. Dept. of For. and Rural Develop., For. Branch., Ottawa. Dept. Pub., 1196: 13pp.
- Kimmins, J.P. 1973. Nutrient Removal Associated with Whole-Tree Logging on Two Different Sites in the Prince George Forest District. Report Submitted to B.C.F.S. Productivity Committee. 100pp. Unpublished.
- King, W.W. and R.L. Schnell. 1972. Biomass estimates of black oak tree components. Tennessee Valley Authority, Norris, Tennessee, Div. For., Fish., and Wildlife Develop., Tech. Note B.1: 24pp.
- Kleinschmit, J. and W. Knigge. 1967. Durch Umwelt und Erbanlagen bedingte Variation der Trockensubstanzerzeugung, Sturktur und Rohdichte junger Fichten (*Picea abies* Karst). (The variation of dry matter production, structure, and density of young spruces (*Picea abies* Karst) as affected by the environment and genotype.) Allg. Forst- u. Jagd Zeitung 138(9): 189-198.
- Krumlik, J.G. 1974. Biomass and Nutrient Distribution in Two Old Growth Forest Ecosystems in South Coastal British Columbia. MSc. Thesis, University of British Columbia, Vancouver, B.C., Faculty of Forestry, 180 pp.
- Krumlik, J.G. and J.P. Kimmins. 1973. Studies of Biomass Distribution and Tree Form in Old Virgin Forests in the Mountains of South Coastal British Columbia. In Anon. 1973, IUFRO Biomass Studies: 363-374.
- Kurucz, J. 1969. Component weights of Douglas-fir, western hemlock and western red cedar biomass for simulation of amount and distribution of forest fuels. MF Thesis, The Univ. of British Columbia, Vancouver, B.C. Faculty of Forestry: 116pp.
- Laundrie, J.F. and J.G. Berbee. 1972. High yields of kraft pulp from rapid-growth hybrid poplar trees. U.S.D.A., For. Serv., For. Prod. Lab., Madison, Wisconsin, Res. Paper FPL 186: 23pp.
- Lieth, H. 1962. Die Stoffproduktion der Pflanzendecke. (The matter production of the vegetation cover.) Vortraege und Diskussionsergebnisse des internationalen oekologischen Symposiums in Stuttgart-Hohenheim vom 4-7 Mai 1960. Fischer, Stuttgart. 156pp.
- Loomis, R.M., R.E. Phares and J.S. Crosby. 1966. Estimating foliage and branchwood quantities in shortleaf pine. For. Sci. 12 (1): 30-39.
- Manwiller, F.G. 1972. Wood and bark properties of spruce pine. U.S.D.A., For. Serv., Southern For. Exp. Sta., New Orleans, La., Res. Paper SO-78: 25pp.
- Marie-Victorin, Frere, edited E. Rouleau. 1964. Flore Laurientienne. Les Presses de L'Universite de Montreal. 925pp.
- Markstrom, D.C. and V.P. Yerkes. 1972. Specific gravity variation with height in Black Hills ponderosa pine. U.S.D.A., For. Serv., Rocky Mountain For. and Range Exp. Sta., Res. Note RM-213: 4pp.
- Martin, W.H. and H. Simard. 1959. Weight as a basis for wood measurement. Canadian International Paper Co., Montreal, Que., Woodlands Sec. Index. No. 1844 (B-6): 149-3 to 9-155 and 168-221.
- Milner, C. and R.E. Hughes. 1968. Methods for the measurement of the land. IBP Handbook 6. Blackwell Scientific Publications, Oxford and Edinburgh: 70pp.

- Moore, J.E. 1971. Forum on "puckerbrush" utilization. Amer. Pulpwood Assoc., 605 Third Ave., New York, N.Y., 10016. Tech. Release 71-R-59: 5pp.
- Morrison, I.K. 1972. Variation with crown position and leaf age in content of seven elements in leaves of Pinus banksiana Lamb. Can. J. For. Res., 2(2): 89-94.
- Newbould, P.J. 1967. Methods for estimating the primary production of forests. IBP handbook No. 2, Blackwell Sci. Pub., Oxford and Edinburgh: 62pp.
- Ovington, J.D. 1957. Dry matter production by Pinus sylvestris L. Annals of Botany, N.S. XXL (82): 287-314.
- Ovington, J.D. 1958. Some biological considerations of forest production. Reprint from the biological productivity of Britain. Pub. by the Inst. of Biology, 41 Queen's Gate, London S.W.7: 8pp.
- Peterson, E.B., Y.H. Chan and J.B. Cragg. 1970. Above-ground standing crop, leaf area, and caloric value in an aspen clone near Calgary, Alberta. Can. For. Serv. Calgary, Alberta. Can. J. Bot. 48: 1459-1469.
- Peck, M.E. 1941. A Manual of the Higher Plants of Oregon. Binford & Mort, Portland, Oregon. 868 pp.
- Post, L.J. 1970. Dry-matter production of mountain maple and balsam fir in northwestern New Brunswick. Ecology 51(3): 548-550.
- Reid, R.W. 1976. Utilization of Aspen for Animal Fodder. Northern Forest Research Centre, Edmonton, Alberta.
- Ribe, J.H. 1973. Puckerbrush weight tables. Life Sci. and Agri. Exp. Sta., Univ. of Maine at Orono. Misc. Rep. 152: 92pp.
- Ribe, J.H. 1973a. A study of multi-stage and dimensional analysis sampling of puckerbrush stands. School of For. Resources, Univ. of Maine at Orono, Working group on forest biomass S4.01-1: 10pp.
- Riedacker, A. 1968. Méthodes indirectes d'estimation de la biomasse des arbres et des peuplements forestiers. (Indirect methods of estimating the biomass of trees and forest publications.) Nat. Inst. of Agr. Res. Nat. Centre of For. Res., Nancy, France: 27pp.
- Riedacker, A. 1971. Measurements of different components of trees for physiological investigations. Ps. 223-240 in Growth and Yield, IUFRO, Univ. of Florida.
- Row, C. and C. Fasick. 1966. Weight-scaling tables by electronic computer. For. Prod. J. 16(8): 41-45.
- Row, C. and S. Guttenberg. 1966. Determining weight-volume relationships for saw logs. For. Prod. J. 16(5): 39-47.
- Schlaegel, B.E. 1973. Estimating the weight yield of Minnesota quaking aspen. (Populus tremuloides Michx.) U.S.D.A., For. Serv., Northern Central For. Exp. Sta., Grand Rapids, Minnesota: 10pp.
- Schmidt, F.L. and D.S. Debell. 1973. Wood production and kraft pulping of short-rotation hardwoods in the Pacific Northwest. Crown Zellerbach, Central Research, Camas, Wash., unpubl. manuscript. 8pp.
- Sharpe, D.M. and W.C. Johnson. 1973. Biomass dynamics of the commercial forests of the Tennessee valley. US-IBP, Eastern Deciduous Forest Biome. rep. 73-23: 13pp.
- Smith, J.H.G. 1972. Tree size and yields in Juvenile Red Alder Stands. Prepared for meeting of the Forestry Section, Northwest Scientific Association, 1972: 35pp.
- Smith, J.H.G. and A. Kozak. 1971. Thickness, moisture content, and specific gravity of inner and outer bark of some Pacific Northwest trees. For. Prod. J. 21(2): 38-40.
- Smith, N.J. 1977. Estimates of Above-Ground Biomass, Net Primary Production and Energy Flows in 8 to 10 Year Old Red Alder (Alnus rubra Bong.) Ecosystems. MF Thesis, University of British Columbia, Vancouver, B.C., Faculty of Forestry: 151pp.
- Steinhilb, H.M. and R.J. Erickson. 1972. Weights and centers of gravity for quaking aspen trees and boles. U.S.D.A. For. Serv., N. Centr. For. Exp. Sta., St. Paul, Minn. Res. Note NC-91: 4pp.
- Steinhilb, H.M. and R.J. Erickson. 1972. Weights and centers of gravity for red pine, white spruce, balsam fir. U.S.D.A. For. Serv., N. Central For. Exp. Sta., St. Paul Minn. Res. Paper NC-75: 7pp.
- Telfer, E.S. 1969. Weight-diameter relationships for 22 woody plant species. Can. J. Bot. 47: 1851-1855.
- Telfer, E.S. 1972. Understory biomass in five forest types in southwestern Nova Scotia. Can. J. Bot. 50: 1263-1267.
- Whittaker, R.H., F.H. Bormann, G.E. Likens and T.G. Siccama. 1974. The Hubbard Brook Ecosystem Study: Forest Biomass and Production. Ecol. Monographs 44: pp. 233-252.
- Whittaker, R.H. and G.M. Woodwell. 1967. Surface area relations of woody plants and forest communities. Amer. J. Bot. 54(8): 931-939.
- Whittaker, R.H. and G.M. Woodwell. 1968. Dimension and production relations of trees and shrubs in the Brookhaven Forest, New York. J. Ecology 56(1): 1-25.

- Whittaker, R.H. and G.M. Woodwell. 1971. Measurement of net primary production of forests. p. 159-184. In Anonymous 1971b.
- Young, H.E., L. Strand and R. Altenberger. 1964. Preliminary fresh and dry weight tables for seven tree species in Maine. Maine Agr. Exp. Sta., Orono, Maine. Tech. Bull. 12: 76pp.
- Young, H.E. 1965. Pound wise and penny foolish. Amer. Pulpwood Assoc. Annual meeting, February 1965, New York, 9pp unpublished.
- Young, H.E. 1966. World forestry based on the complete tree concept. Sixth World Forestry Congress, Madrid, Spain: 5pp.
- Young, H.E. 1968. Challenge of complete tree utilization. For. Prod. J. 18(4): 83-86.
- Young, H.E. 1968. Quantum increases in fiber production. Amer. Soc. of Agr. Engineers. Proc. For. Eng. Conf., Saint Joseph, Mich., ps. 102-103, and 105.
- Young, H.E. 1971. Preliminary estimates of bark percentages and chemical elements in complete trees of eight species in Maine. For. Prod. J. 21(5): 56-59.
- Young, H.E. 1972. Puckerbrush utilization for paper products. School of For. Res., Univ. of Maine at Orono. Paper prepared for annual meeting, Tappi, New York, N.Y., February 14-16, 1972: 12pp unpublished.
- Young, H.E. 1975. The enormous potential of the forests: a positive rebuttal to Grantham and Ellis. J. Forestry 73: 99-102.
- Young, H.E. 1976. Muka: a good Russian Idea. J. of For. 74(3): 1pp.
- Young, H.E. and P.M. Carpenter. 1967. Weight, nutrient element and productivity studies of seedlings and saplings of eight tree species in natural ecosystems. Maine Agr. Exp. Sta., Univ. of Maine at Orono. Tech. Bull 28: 39pp.
- Young, H.E., P. Carpenter and R. Altenberger. 1965. Preliminary tables of some chemical elements in seven tree species in Maine., Maine Agr. Exp. Sta., Orono, Maine. Tech. Bull. 20: 88pp.
- Young, H.E. and A.J. Chase. 1965. Fiber weight and pulping characteristics of the logging residue of seven tree species in Maine. Maine Agr. Exp. Sta., Univ. of Maine at Orono. Tech. Bull. 17: 43pp.
- Young, H.E., C.B. Gammon and M. Ashley. 1964. Potential fiber from balsam fir, white pine, hemlock, white birch, and aspen logging residue. Tappi 47(9): 555-557.
- Young, H.E., C.B. Gammon and L.E. Hoar. Potential fiber from red spruce and red maple logging residue. Tappi 46(4): 256-259.
- Young, H.E. and V.P. Guinn. 1966. Chemical elements in complete mature trees of seven species in Maine. Tappi 49(5): 190-197.
- Young, H.E., L. Hoar and M. Ashley. 1965. Weight of wood substance for components of seven tree species. Univ. of Maine at Orono, Maine. Tappi 48(8): 466-469.
- Young, H.E., L. Strand and R. Altenberger. 1964. Preliminary fresh and dry weight tables for seven tree species in Maine. Maine Agr. Exp. Sta., Orono, Maine. Tech. Bull. 12: 76pp.
- Zyryaev, A.G. 1964. The relationship between needle mass and current increment in Larix Sibirica. Can. Dept. of For. Translation from Lesnoye Khozaystvo, 17(4): 28-29.
- Zavitkovski, J. and R.D. Stevens. 1972. Primary Productivity of Red Alder Ecosystems. In Anon. 1972, Ecology: 235-241.

Appendix Regression equations for biomass of individual species
(for explanation of details see text)

SPECIES (Reference) Equation Title	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Alder</u> Speckled (Ribe, 1973)					
Foliage	FW-g	$\log Y = 2.5987 + 1.2510 \log D_2$	30	.70	2.5-7.5 (cm)
Branches	FW-g	$\log Y = 2.7096 + 1.4877 \log D_2$	30	.77	2.5-7.5 (cm)
Stem	FW-g	$\log Y = 3.0095 + 2.4534 \log D_2$	30	.93	2.5-7.5 (cm)
Foliage	ODW-g	$\log Y = 2.1759 + 1.2463 \log D_2$	30	.67	2.5-7.5 (cm)
Branches	ODW-g	$\log Y = 2.4072 + 1.5184 \log D_2$	30	.78	2.5-7.5 (cm)
Stem	ODW-g	$\log Y = 2.7051 + 2.5847 \log D_2$	30	.93	2.5-7.5 (cm)
<u>Alder</u> Red (Smith, J., 1972)					
Tree	FW-g	$Y = 1942.91 + 93.4572 D_6$	230	.86	3-149 (mm)
Tree	FW-g	$Y = 933.341 + 54.737 (D_9)^2 H_1$	230	.90	3-149 (mm)
Foliage	FW-g	$Y = 454.086 + 53578 BA$	230	.52	3-149 (mm)
Stem	FW-g	$Y = 933.984 + 277680 BA$	230	.78	3-149 (mm)
Branches	FW-g	$Y = 251.325 + 98022.2 BA$	230	.70	3-149 (mm)
Tree	FW-g	$Y = 1634.34 + 428724 BA$	230	.81	3-149 (mm)
Tree	FW-g	$\log Y = 3.67006 + 2.03887 \log D_{11} - 1.64423 \log H_1$	230	.94	3-149 (mm)
Tree	FW-g	$\log Y = 3.47024 + 1.94582 \log D_{11}$	230	.94	3-149 (mm)
<u>Alder</u> Red (Smith, N., 1977)					
Stem, wood	ODW-kg	$Y = 1.31625 + .346084 ((D_5)^2 H_3)/100 + .573351 D_9/100$	72	.97	4.2-6.0 (cm)
Stem, bark	ODW-kg	$Y = .370339 + .0227208 ((D_5)^2 H_3)/100 + .140593 D_9/100$	72	.95	4.2-6.0 (cm)
Stem, wood & bark	ODW-kg	$Y = 1.686589 + .368048 ((D_5)^2 H_3)/100 + .713944 D_9/100$	72	.98	4.2-6.0 (cm)
Total Crown (including reproductive organs)	ODW-kg	$Y = .940575 + .566637 D_9/100 + .0484421 CL_m/CW_m$	66	.69	4.2-6.0 (cm)
Branches <.6 cm and foliage	ODW-kg	$Y = .429308 + .422831 D_9/100 + .0487709 CL_m/CW_m$	66	.68	4.2-6.0 (cm)
Branches <.6 cm	ODW-kg	$Y = .487593 + .130159 D_9/100 + .01159 CL_m/CW_m$	66	.64	4.2-6.0 (cm)

Appendix

SPECIES (Reference)	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Alder</u>					
<u>Red</u>					
(Zavitkovski & Stevens, 1972)					
Crowns	ODW-kg	$Y = .01 + .48 IV - .0009 (IV)^2$	91	.75	1.25 - 152.5 (cm) (l.b.)
Stems	ODW-kg	$Y = .02 + 1.60 IV - .0005 (IV)^2$	91	.99	1.25 - 152.5 (cm)
Tree	ODW-kg	$Y = .02 + 2.09 IV - .0015 (IV)^2$	119	.98	1.25 - 152.5 (cm)
Roots (estimated)	ODW-kg	$Y = .1 + .48 IV - .0005 (IV)^2$	-	-	1.25 - 152.5 (cm)
Tops	ODW-MT/ha	$Y = 17.15 + 5.24 CH + .143 (CH)^2$	50	.98	1.25 - 152.5 (cm)
<u>Aspen</u>					
<u>sp.</u>					
(Young et al., 1964)					
Tree tot.	FW-lbs	$\ln Y = .6800 + 2.2234 \ln D_1 + .3390 \ln H_1$	14	.97	14.5 - 24.5 (cm)
Tree tot.	ODW-lbs.	34% of FW from above equation	14	.97	14.5 - 24.5 (cm)
Tree tot. - Roots < 4"	FW-lbs	$\ln Y = -1.5880 + 2.0013 \ln D_1 + 1.0100 \ln H_1$	14	.97	14.5 - 24.5 (cm)
Tree tot. - Roots < 4"	ODW-lbs	35% of FW from above equation	14	.97	14.5 - 24.5 (cm)
Tree a.s. 6"	FW-lbs	$\ln Y = 2.066 + 2.1679 \ln D_1 + .4292 \ln H_1$	14	.97	14.5 - 24.5 (cm)
Tree a.s. 6"	ODW-lbs	35% of FW from above equation	14	.97	14.5 - 24.5 (cm)
Stem a.s. 6"	FW-lbs	$\ln Y = -1.4518 + 1.8801 \ln D_1 + .9332 \ln H_1$	14	.98	14.5 - 24.5 (cm)
Stem a.s. 6"	ODW-lbs	38% of FW from above equation	14	.98	14.5 - 24.5 (cm)
Bole a.s. 6", t. 4"	FW-lbs	$\ln Y = -1.9730 + 2.0215 \ln D_1 + .9670 \ln H_1$	14	.97	14.5 - 24.5 (cm)
Bole a.s. 6", t. 4"	ODW-lbs	38% of FW from above equation	14	.97	14.5 - 24.5 (cm)
All branches	FW-lbs	$\ln Y = 2.5720 + 3.7019 \ln D_1 - 1.3520 \ln H_1$	14	.88	14.5 - 24.5 (cm)
All branches	ODW-lbs	21% of FW from above equation	14	.88	14.5 - 24.5 (cm)
Branches < 1 in	FW-lbs	$\ln Y = 5.0497 + 2.8069 \ln D_1 - 1.6230 \ln H_1$	14	.74	14.5 - 24.5 (cm)
Branches < 1 in	ODW-lbs	9% of FW from above equation	14	.74	14.5 - 24.5 (cm)
Stump 6" & roots > 1 in	FW-lbs	$\ln Y = -11.6867 + 1.6151 \ln D_1 + 3.2358 \ln H_1$	14	.78	14.5 - 24.5 (cm)
Stump 6" & roots > 1 in	ODW-lbs	34% of FW from above equation	14	.78	14.5 - 24.5 (cm)
Roots > 1 in	FW-lbs	$\ln Y = -15.5139 + 1.2752 \ln D_1 + 4.2545 \ln H_1$	14	.64	14.5 - 24.5 (cm)
Roots > 1 in	ODW-lbs	30% of FW from above equation	14	.64	14.5 - 24.5 (cm)
Roots < 1 in	FW-lbs	$\ln Y = -1.6126 + 1.6306 \ln D_1 + .3564 \ln H_1$	14	.08	14.5 - 24.5 (cm)
Roots < 1 in	ODW-lbs	19% of FW from above equation	14	.08	14.5 - 24.5 (cm)

Appendix

SPECIES (Reference Equation Title	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Aspen</u> sp. (Young and Carpenter, 1967)					
Foliage	FW-g	$\ln Y = -0.228 + 2.328 \ln H_1$	6	.93	0.0 - 75 (mm)
Branches	FW-g	$\ln Y = -2.613 + 3.130 \ln H_1$	6	.89	0.0 - 75 (mm)
Stem	FW-g	$\ln Y = -1.482 + 3.364 \ln H_1$	6	.99	0.0 - 75 (mm)
Roots	FW-g	$\ln Y = -0.388 + 2.707 \ln H_1$	6	.93	0.0 - 75 (mm)
Tree tot.	FW-g	$\ln Y = 0.167 + 3.034 \ln H_1$	6	.97	0.0 - 75 (mm)
Foliage	ODW-g	$\ln Y = -1.118 + 2.352 \ln H_1$	6	.95	0.0 - 75 (mm)
Branches	ODW-g	$\ln Y = -3.225 + 3.144 \ln H_1$	6	.92	0.0 - 75 (mm)
Stem	ODW-g	$\ln Y = -2.064 + 3.334 \ln H_1$	6	.99	0.0 - 75 (mm)
Roots	ODW-g	$\ln Y = -1.331 + 2.754 \ln H_1$	6	.94	0.0 - 75 (mm)
Tree tot	ODW-g	$\ln Y = -0.653 + 3.067 \ln H_1$	6	.97	0.0 - 75 (mm)
<u>Aspen</u> Trembling (Quaking) (Rbe, 1973)					
Foliage	FW-g	$\log Y = 2.4385 + 1.6566 \log D_2$	30	.76	2.5 - 15.0 (cm)
Branches	FW-g	$\log Y = 2.5550 + 1.7990 \log D_2$	30	.85	2.5 - 15.0 (cm)
Stem	FW-g	$\log Y = 3.0207 + 2.6528 \log D_2$	30	.99	2.5 - 15.0 (cm)
Foliage	ODW-g	$\log Y = 2.0243 + 1.6796 \log D_2$	30	.75	2.5 - 15.0 (cm)
Branches	ODW-g	$\log Y = 2.2178 + 1.8545 \log D_2$	30	.85	2.5 - 15.0 (cm)
Stem	ODW-g	$\log Y = 2.6672 + 2.7859 \log D_2$	30	.99	2.5 - 15.0 (cm)
<u>Aspen</u> Trembling (Quaking) (Peterson, Chan and Cragg, 1970)					
Tree	ODW-kg	$\log Y = -1.1115 + 2.3466 \log D_2$	49	.98 (r)	4.5 - 33.0 (cm)
Stem	ODW-kg	$\log Y = -1.1893 + 2.3564 \log D_2$	49	.97 (r)	4.5 - 33.0 (cm)
Branches	ODW-kg	$\log Y = -2.0978 + 2.3708 \log D_2$	49	.97 (r)	4.5 - 33.0 (cm)
Foliage	ODW-kg	$\log Y = -2.3011 + 1.9742 \log D_2$	49	.97 (r)	4.5 - 33.0 (cm)
Crown	ODW-kg	$\log Y = -1.9282 + 2.2990 \log D_2$	49	.97 (r)	4.5 - 33.0 (cm)

Appendix

SPECIES (Reference)	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Aspen</u>					
Trembling (Quaking) (Peterson, Chan and Cragg, 1970)					
.... cont'd					
Tree	ODW-kg	$\log Y = -3.2198 + 0.8993 \log D_2$	49	.99 (r)	4.5 - 33.0 (cm)
Stem	ODW-kg	$\log Y = -3.3211 + 0.9061 \log D_8$	49	.99 (r)	4.5 - 33.0 (cm)
Branches	ODW-kg	$\log Y = -4.1237 + 0.8870 \log D_8$	49	.95 (r)	4.5 - 33.0 (cm)
Foliage	ODW-kg	$\log Y = -4.0324 + 0.7478 \log D_8$	49	.96 (r)	4.5 - 33.0 (cm)
Crown	ODW-kg	$\log Y = -3.8965 + 0.8610 \log D_8$	49	.95 (r)	4.5 - 33.0 (cm)
Tree	ODW-kg	$\log Y = -8.1917 + 3.1855 \log D_4$	49	.92 (r)	4.5 - 33.0 (cm)
Stem	ODW-kg	$\log Y = -8.4267 + 3.2424 \log D_4$	49	.93 (r)	4.5 - 33.0 (cm)
Branches	ODW-kg	$\log Y = -8.3477 + 2.9097 \log D_4$	49	.82 (r)	4.5 - 33.0 (cm)
Foliage	ODW-kg	$\log Y = -7.8891 + 2.5541 \log H_2$	49	.87 (r)	4.5 - 33.0 (cm)
Crown	ODW-kg	$\log Y = -8.0216 + 2.8328 \log H_2$	49	.83 (r)	4.5 - 33.0 (cm)
Branches	ODW-kg	$\log Y = -1.3967 + 2.1979 \log D$	49	.98 (r)	4.5 - 33.0 (cm)
Foliage	ODW-kg	$\log Y = -1.7121 + 1.8229 \log D$	49	.98 (r)	4.5 - 33.0 (cm)
Crown	ODW-kg	$\log Y = -1.2500 + 2.1336 \log D$	49	.99 (r)	4.5 - 33.0 (cm)
Branches	ODW-kg	$\log Y = -7.1668 + 2.9132 \log CL$	49	.86 (r)	4.5 - 33.0 (cm)
Foliage	ODW-kg	$\log Y = -6.6624 + 2.4816 \log CL$	49	.88 (r)	4.5 - 33.0 (cm)
Crown	ODW-kg	$\log Y = -6.8470 + 2.8263 \log CL$	49	.87 (r)	4.5 - 33.0 (cm)
Branches	ODW-kg	$\log Y = -0.8411 + 0.6348 \log CR$	49	.12 (r)	4.5 - 33.0 (cm)
Foliage	ODW-kg	$\log Y = -1.0023 + 0.3701 \log CR$	49	.08 (r)	4.5 - 33.0 (cm)
Crown	ODW-kg	$\log Y = -0.6744 + 0.5936 \log CR$	49	.12 (r)	4.5 - 33.0 (cm)
Stem & Branches	ODW-kg	$\log Y = -1.1344 + 2.3567 \log D_2$	49	.98 (r)	4.5 - 33.0 (cm)
Stem & Branches	ODW-kg	$\log Y = -3.2531 + 0.9035 \log D_8$	49	.99 (r)	4.5 - 33.0 (cm)
Stem & Branches	ODW-kg	$\log Y = -8.2569 + 3.2033 \log H_2$	49	.92 (r)	4.5 - 33.0 (cm)
Branches	ODW-kg	$Y = 0.0639 D^{2.0291}$	49	.99 (r)	4.5 - 33.0 (cm)
Foliage	ODW-kg	$Y = 0.0301 D^{1.6238}$	49	.99 (r)	4.5 - 33.0 (cm)

Appendix

SPECIES (Reference) Equation Title	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Aspen</u>					
<u>Trenbiling (Quaking)</u> (Telfer, 1969)					
Tree	ODW-g	$\ln Y = -2.920 + 2.715 \ln D_4$	20	.99	2.3 - 40.6 (cm)
Foliage	ODW-g	$\ln Y = -2.677 + 2.156 \ln D_4$	20	.96	2.3 - 40.6 (cm)
<u>Bayberry</u> (Telfer, 1969)					
Tree	ODW-g	$\ln Y = -2.494 + 2.867 \ln D_4$	20	.98	1.9 - 12.0 (mm)
Foliage	ODW-g	$\ln Y = -2.913 + 2.529 \ln D_4$	20	.93	1.9 - 12.0 (mm)
<u>Beech</u> (Telfer, 1969)					
Tree	ODW-g	$\ln Y = -3.647 + 2.906 \ln D_4$	20	.99	1.8 - 35.5 (mm)
Foliage	ODW-g	$\ln Y = -3.910 + 2.354 \ln D_4$	20	.99	1.8 - 35.5 (mm)
<u>Beech</u> (Ribe, 1973)					
Foliage	FW-g	$\log Y = 2.4271 + 1.7292 \log D_2$	19	.85	2.5 - 15.0 (cm)
Branches	FW-g	$\log Y = 2.8179 + 1.5359 \log D_2$	19	.80	2.5 - 15.0 (cm)
Stem	FW-g	$\log Y = 3.2794 + 2.4751 \log D_2$	19	.99	2.5 - 15.0 (cm)
Foliage	ODW-g	$\log Y = 2.0660 + 1.8089 \log D_2$	19	.85	2.5 - 15.0 (cm)
Branches	ODW-g	$\log Y = 2.5983 + 1.5402 \log D_2$	19	.79	2.5 - 15.0 (cm)
Stem	ODW-g	$\log Y = 3.0692 + 2.4868 \log D_2$	19	.99	2.5 - 15.0 (cm)
<u>Birch</u> Gray (Ribe, 1973)					
Foliage	FW-g	$\log Y = 2.3679 + 1.5698 \log D_2$	30	.86	2.5 - 15.0 (cm)
Branches	FW-g	$\log Y = 2.7349 + 1.6033 \log D_2$	30	.84	2.5 - 15.0 (cm)
Stem	FW-g	$\log Y = 3.2137 + 2.4848 \log D_2$	30	.99	2.5 - 15.0 (cm)
Foliage	ODW-g	$\log Y = 1.8730 + 1.6376 \log D_2$	30	.87	2.5 - 15.0 (cm)
Branches	ODW-g	$\log Y = 2.4831 + 1.6163 \log D_2$	30	.81	2.5 - 15.0 (cm)
Stem	ODW-g	$\log Y = 2.9508 + 2.5139 \log D_2$	30	.99	2.5 - 15.0 (cm)

Appendix

SPECIES (Reference)	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Birch</u> White (paper) (Baskerville, 1965)					
Tree	FW-lbs	$\log Y = 0.736 + 2.17 \log D_1$	24	.95	3 - 9 (cm)
Tree	ODW-lbs	$\log Y = 0.236 + 2.48 \log D_1$	24	.98	3 - 9 (cm)
Stem wood	ODW-lbs	$\log Y = 0.312 + 2.36 \log D_1$	24	.97	3 - 9 (cm)
Stem bark	ODW-lbs	$\log_{100} Y = 1.32 + 2.35 \log D_1$	24	.90	3 - 9 (cm)
Branches	ODW-lbs	$\log Y = -1.006 + 3.30 \log D_1$	24	.86	3 - 9 (cm)
Foliage	ODW-lbs	$\log_{100} Y = 0.730 + 2.94 \log D_1$	25	.90	3 - 9 (cm)
Dead Branches	ODW-lbs	$\log_{100} Y = 0.679 + 3.30 \log D_1$	20	.68	3 - 9 (cm)
<u>Birch</u> White (paper) (Ribe, 1973)					
Foliage	FW-g	$\log Y = 2.6161 + 1.6161 \log D_2$	30	.89	2.5 - 15.0 (cm)
Branches	FW-g	$\log Y = 2.6961 + 1.6886 \log D_2$	30	.88	2.5 - 15.0 (cm)
Stem	FW-g	$\log Y = 2.7167 + 3.2104 \log D_2$	30	.99	2.5 - 15.0 (cm)
Foliage	ODW-g	$\log Y = 2.1587 + 1.7020 \log D_2$	30	.89	2.5 - 15.0 (cm)
Branches	ODW-g	$\log Y = 2.4059 + 1.7304 \log D_2$	30	.88	2.5 - 15.0 (cm)
Stem	ODW-g	$\log Y = 2.4481 + 3.2640 \log D_2$	30	.99	2.5 - 15.0 (cm)
<u>Birch</u> White (paper) (Young et al. 1964)					
Tree tot.	FW-lbs	$\ln Y = .8025 + 2.2234 \ln D_1 + .3390 \ln H_1$	17	.97	15.3 - 29.0 (cm)
Tree tot.	ODW-lbs	42% of FW from above equation	17	.97	15.3 - 29.0 (cm)
Tree tot. - roots < 4"	FW-lbs	$\ln Y = .6040 + 2.2236 \ln D_1 + .3635 \ln H_1$	17	.98	15.3 - 29.0 (cm)
Tree tot. - roots < 4"	ODW-lbs	43% of FW from above equation	17	.98	15.3 - 29.0 (cm)
Tree a.s. 6"	FW-lbs	$\ln Y = .3062 + 2.1679 \ln D_1 + .4292 \ln H_1$	17	.97	15.3 - 29.0 (cm)
Tree a.s. 6"	ODW-lbs	42% of FW from above equation	17	.97	15.3 - 29.0 (cm)
Stem a.s. 6"	FW-lbs	$\ln Y = -1.3721 + 1.8801 \ln D_1 + .9332 \ln H_1$	17	.98	15.3 - 29.0 (cm)
Stem a.s. 6"	ODW-lbs	43% of FW from above equation	17	.98	15.3 - 29.0 (cm)

Appendix

SPECIES (Reference) Equation Title	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Birch</u>					
White (paper) (Young et al. 1964)					
... cont'd					
Bole a.s. 6", t 4"	FW-lbs	$\ln Y = -1.8959 + 2.0215 \ln D_1 + .9670 \ln H_1$	17	.98	15.3 - 29.0 (cm)
Bole a.s. 6", t 4"	ODW-lbs	48% of FW from above equation	17	.98	15.3 - 29.0 (cm)
All branches	FW-lbs	$\ln Y = 8.6990 + 5.1205 \ln D_1 - 3.5493 \ln H_1$	17	.69	15.3 - 29.0 (cm)
All branches	ODW-lbs	20% of FW from above equation	17	.69	15.3 - 29.0 (cm)
Branches < 1"	FW-lbs	$\ln Y = 5.5076 + 3.5204 \ln D_1 - 2.0449 \ln H_1$	17	.61	15.3 - 29.0 (cm)
Branches < 1"	ODW-lbs	13% of FW from above equation	17	.61	15.3 - 29.0 (cm)
Stump 6" & roots > 1"	FW-lbs	$\ln Y = 3.8968 + 3.0243 \ln D_1 - 1.2598 \ln H_1$	17	.76	15.3 - 29.0 (cm)
Stump 6" & roots > 1"	ODW-lbs	48% of FW from above equation	17	.76	15.3 - 29.0 (cm)
Roots > 1"	FW-lbs	$\ln Y = 5.5082 + 3.2590 \ln D_1 - 1.8989 \ln H_1$	17	.65	15.3 - 29.0 (cm)
Roots > 1"	ODW-lbs	44% of FW from above equation	17	.65	15.3 - 29.0 (cm)
Roots < 1"	FW-lbs	$\ln Y = -1.2539 + 1.6306 \ln D_1 + .3564 \ln H_1$	17	.46	15.3 - 29.0 (cm)
Roots < 1"	ODW-lbs	25% of FW from above equation	17	.46	15.3 - 29.0 (cm)
<u>Birch</u>					
White (paper) (Young and Carpenter, 1967)					
Foliage	FW-g	$\ln Y = 1.310 + 2.004 \ln H_1$	10	.97	0 - 75 (mm)
Branches	FW-g	$\ln Y = 0.311 + 2.312 \ln H_1$	10	.97	0 - 75 (mm)
Stem	FW-g	$\ln Y = -0.066 + 2.892 \ln H_1$	10	.99	0 - 75 (mm)
Roots	FW-g	$\ln Y = 0.181 + 2.431 \ln H_1$	10	.97	0 - 75 (mm)
Tree tot.	FW-g	$\ln Y = 1.900 + 2.473 \ln H_1$	10	.98	0 - 75 (mm)
Foliage	ODW-g	$\ln Y = -0.007 + 2.116 \ln H_1$	10	.97	0 - 75 (mm)
Branches	ODW-g	$\ln Y = -0.157 + 2.276 \ln H_1$	10	.97	0 - 75 (mm)
Stem	ODW-g	$\ln Y = -0.212 + 2.739 \ln H_1$	10	.98	0 - 75 (mm)
Roots	ODW-g	$\ln Y = -0.263 + 2.375 \ln H_1$	10	.97	0 - 75 (mm)
Tree tot.	ODW-g	$\ln Y = 1.183 + 2.492 \ln H_1$	10	.98	0 - 75 (mm)

Appendix

SPECIES (Reference)	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Birch</u>					
Yellow (Ribe, 1973)					
Foliage	FW-g	$\log Y = 2.4488 + 1.8961 \log D_2$	30	.90	2.5 — 15.0 (cm)
Branches	FW-g	$\log Y = 2.7713 + 1.6306 \log D_2$	30	.90	2.5 — 15.0 (cm)
Stem	FW-g	$\log Y = 3.1837 + 2.5206 \log D_2$	30	.99	2.5 — 15.0 (cm)
Foliage	ODW-g	$\log Y = 1.9962 + 1.9683 \log D_2$	30	.90	2.5 — 15.0 (cm)
Branches	ODW-g	$\log Y = 2.5345 + 1.6179 \log D_2$	30	.89	2.5 — 15.0 (cm)
Stem	ODW-g	$\log Y = 2.9670 + 2.5330 \log D_2$	30	.99	2.5 — 15.0 (cm)
<u>Blueberry</u>					
Lowbush (Telfer, 1969)					
Tree	ODW-g	$\ln Y = -3.978 + 3.706 \ln D_4$	20	.94	1.4 — 5.6 (mm)
Foliage	ODW-g	$\ln Y = -4.404 + 3.034 \ln D_4$	20	.74	1.4 — 5.6 (mm)
<u>Blueberry</u>					
Lowbush (Whittaker and Woodwell, 1968)					
Stem	ODW-g	$\log Y = 1.2758 + 2.9605 \log D_5$	15	.85 (r)	No data available
Stem wood	ODW-g	$\log Y = 1.1773 + 3.1513 \log D_5$	15	.83 (r)	No data available
Stem bark	ODW-g	$\log Y = 0.7705 + 2.8065 \log D_5$	15	.81 (r)	No data available
Branches	ODW-g	$\log Y = 1.1999 + 2.6115 \log D_5$	15	.86 (r)	No data available
Tree	ODW-g	$\log Y = 1.6937 + 2.4995 \log D_5$	15	.87 (r)	No data available
<u>Cedar</u>					
Northern White (large) (Dyer, 1967)					
Tree tot.	FW-lbs	$\log Y = -2.07 + 1.59 \log D_1 + 1.29 \log H_1$	21	.90	14.0 (cm)
Tree tot. — Roots < 4"	FW-lbs	$\log Y = -2.04 + 1.62 \log D_1 + 1.25 \log H_1$	21	.91	14.0 (cm)
Tree	FW-lbs	$\log Y = -2.63 + 1.53 \log D_1 + 1.41 \log H_1$	21	.90	14.0 (cm)
Stem	FW-lbs	$\log Y = -3.94 + 1.46 \log D_1 + 1.71 \log H_1$	21	.93	14.0 (cm)
Bole, s ₂ , t ₇	FW-lbs	$\log Y = -5.44 + 1.56 \log D_1 + 2.01 \log H_1$	21	.95	14.0 (cm)

Appendix

SPECIES (Reference)	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Cedar</u>					
Northern White (large)					
(Dyer, 1967)					
..... cont'd					
Stump ? & Roots > 1"	FW-lbs	$\log Y = -1.96 + 1.90 \log D_1 + 0.61 \log H_1$	21	.89	14.0 (cm)
Roots > 1"	FW-lbs	$\log Y = -4.39 + 1.78 \log D_1 + 1.18 \log H_1$	21	.77	14.0 (cm)
Roots < 1"	FW-lbs	$\log Y = -7.96 + 1.31 \log D_1 + 2.06 \log H_1$	21	.55	14.0 (cm)
Tree tot.	ODW-lbs	$\log Y = -3.29 + 1.53 \log D_1 + 1.40 \log H_1$	21	.92	14.0 (cm)
Tree tot. — Roots < 4"	ODW-lbs	$\log Y = -3.30 + 1.53 \log D_1 + 1.39 \log H_1$	21	.93	14.0 (cm)
Tree	ODW-lbs	$\log Y = -3.97 + 1.41 \log D_1 + 1.59 \log H_1$	21	.91	14.0 (cm)
Stem	ODW-lbs	$\log Y = -4.62 + 1.29 \log D_1 + 1.79 \log H_1$	21	.92	> 14.0 (cm)
Bole, s?, t?	ODW-lbs	$\log Y = -6.02 + 1.39 \log D_1 + 2.06 \log H_1$	21	.95	> 14.0 (cm)
Stump ? & Roots > 1"	ODW-lbs	$\log Y = -2.81 + 2.00 \log D_1 + 0.54 \log H_1$	21	.90	> 14.0 (cm)
Roots > 1"	ODW-lbs	$\log Y = -5.70 + 1.96 \log D_1 + 1.51 \log H_1$	21	.79	> 14.0 (cm)
Roots < 1"	ODW-lbs	$\log Y = -8.82 + 1.84 \log D_1 + 1.52 \log H_1$	21	.51	> 14.0 (cm)
<u>Cedar</u>					
Northern White (small)					
(Dyer, 1967)					
Foliage	FW-g	$\log Y = 2.58 + 2.03 \log H_1$	36	.94	< 14.0 (cm)
Foliage	ODW-g	$\log Y = 1.59 + 2.07 \log H_1$	36	.94	< 14.0 (cm)
Branches	FW-g	$\log Y = 1.20 + 2.40 \log H_1$	36	.90	< 14.0 (cm)
Branches	ODW-g	$\log Y = 0.59 + 2.37 \log H_1$	36	.89	< 14.0 (cm)
Stem	FW-g	$\log Y = 0.59 + 2.98 \log H_1$	36	.97	< 14.0 (cm)
Stem	ODW-g	$\log Y = -0.32 + 3.07 \log H_1$	36	.95	< 14.0 (cm)
Roots	FW-g	$\log Y = 1.31 + 2.46 \log H_1$	36	.94	< 14.0 (cm)
Roots	ODW-g	$\log Y = 0.60 + 2.40 \log H_1$	36	.93	< 14.0 (cm)
Tree tot.	FW-g	$\log Y = 3.02 + 2.41 \log H_1$	36	.96	< 14.0 (cm)
Tree tot.	ODW-g	$\log Y = 2.21 + 2.42 \log H_1$	36	.96	< 14.0 (cm)

Appendix

SPECIES (Reference)	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Cedar</u>					
Northern White (Young and Carpenter, 1967)					
Foliage	FW-g	$\ln Y = 2.578 + 2.026 \ln H_1$	34	.94	2.5 — 10.0 (cm)
Branches	FW-g	$\ln Y = 1.205 + 2.404 \ln H_1$	34	.90	2.5 — 10.0 (cm)
Stem	FW-g	$\ln Y = 0.595 + 2.985 \ln H_1$	34	.97	2.5 — 10.0 (cm)
Roots	FW-g	$\ln Y = 1.311 + 2.455 \ln H_1$	34	.94	2.5 — 10.0 (cm)
Tree tot.	FW-g	$\ln Y = 3.022 + 2.409 \ln H_1$	34	.96	2.5 — 10.0 (cm)
Foliage	ODW-g	$\ln Y = 1.594 + 2.073 \ln H_1$	34	.95	2.5 — 10.0 (cm)
Branches	ODW-g	$\ln Y = 0.587 + 2.371 \ln H_1$	34	.89	2.5 — 10.0 (cm)
Stem	ODW-g	$\ln Y = -0.321 + 3.074 \ln H_1$	34	.95	2.5 — 10.0 (cm)
Roots	ODW-g	$\ln Y = 0.605 + 2.400 \ln H_1$	34	.93	2.5 — 10.0 (cm)
Tree tot.	ODW-g	$\ln Y = 2.213 + 2.421 \ln H_1$	34	.96	2.5 — 10.0 (cm)
<u>Western Red (Kurucz, 1961)</u>					
Tree	ODW-lbs	$Y = 0.037297 D_6$	113	.90	2.5 — 120 (cm)
Stem - wood	ODW-lbs	$Y = 0.028292 D_6$	113	.90	2.5 — 120 (cm)
Stem - bark	ODW-lbs	$Y = 0.003928 D_6$	113	.93	2.5 — 120 (cm)
Crown	ODW-lbs	$Y = 0.301496 DC_1 + 0.000072 DC_2$	113	.79	2.5 — 120 (cm)
Branches > 3/4 in	ODW-lbs	$Y = 0.099398 DC_1 + 0.000056 DC_2$	113	.77	2.5 — 120 (cm)
Branches 1/4 - 3/4 in	ODW-lbs	$Y = 0.029247 DC_1 + 0.000008 DC_2$	113	.78	2.5 — 120 (cm)
Branches < 1/4 in	ODW-lbs	$Y = 0.022392 DC_1 + 0.000002 DC_2$	113	.82	2.5 — 120 (cm)
Foliage	ODW-lbs	$Y = 0.150460 DC_1 + 0.000006 DC_2$	113	.81	2.5 — 120 (cm)
<u>Cherry</u>					
Choke (Ribe, 1973)					
Foliage	FW-g	$\log Y = 2.4389 + 1.2857 \log D_2$	16	.60	2.5 — 7.5 (cm)
Branches	FW-g	$\log Y = 2.7898 + 1.2164 \log D_2$	16	.57	2.5 — 7.5 (cm)
Stem	FW-g	$\log Y = 3.0997 + 1.9898 \log D_2$	16	.84	2.5 — 7.5 (cm)

Appendix

SPECIES (Reference)	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Cherry</u>					
Choke (Ribe, 1973)					
.... cont'd					
Foliage	ODW-g	$\log Y = 2.0526 + 1.3307 \log D_2$	16	.60	2.5 - 7.5 (cm)
Branches	ODW-g	$\log Y = 2.5540 + 1.2191 \log D_2$	16	.56	2.5 - 7.5 (cm)
Stem	ODW-g	$\log Y = 2.8760 + 2.0038 \log D_2$	16	.85	2.5 - 7.5 (cm)
<u>Cherry</u>					
Pin (Ribe, 1973)					
Foliage	FW-g	$\log Y = 2.5825 + 1.9060 \log D_2$	30	.90	2.5 - 15.0 (cm)
Branches	FW-g	$\log Y = 2.7161 + 1.8263 \log D_2$	30	.88	2.5 - 15.0 (cm)
Stem	FW-g	$\log Y = 3.2025 + 2.1947 \log D_2$	30	.98	2.5 - 15.0 (cm)
Foliage	ODW-g	$\log Y = 2.0974 + 1.9784 \log D_2$	30	.90	2.5 - 15.0 (cm)
Branches	ODW-g	$\log Y = 2.4033 + 1.8755 \log D_2$	30	.87	2.5 - 15.0 (cm)
Stem	ODW-g	$\log Y = 2.9117 + 2.2988 \log D_2$	30	.98	2.5 - 15.0 (cm)
<u>Fern</u>					
Sweet (Telfer, 1969)					
Tree	ODW-g	$\ln Y = -3.831 + 3.314 \ln D_4$	20	.97	2.1 - 14.8 (mm)
Foliage	ODW-g	$\ln Y = -4.331 + 3.014 \ln D_4$	20	.97	2.1 - 14.8 (mm)
<u>Fern</u>					
Bracken (Stanek and State, 1978)					
Tree	ODW-g/m ²	$Y = 1.02 V + .009 V^2$	65	.91	- - -
Tree	ODW-g/m ²	$Y = 10.55 ME$	35	.84	- - -
<u>Fir</u>					
Amabilis (Pacific Silver)					
(Krumlik and Kimmins, 1973)					
Tree, a.s. 30 cm	ODW-kg	$Y = 168.704 + 116.779 D_7$	7	.99	31.0 - 90.4 (cm)
Stem, a.s. 30 cm, wood	ODW-kg	$Y = 80.824 + 90.478 D_7$	7	.99	31.0 - 90.4 (cm)
Stem, a.s. 30 cm, bark	ODW-kg	$Y = -42.324 + 1052.280 BA$	7	.95	31.0 - 90.4 (cm)

Appendix

SPECIES (Reference) Equation Title	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
Fir					
Amabilis (Pacific Silver) (Krumlik and Kimmins, 1973)					
Branches, > 2.5 cm cont'd	ODW-kg	$Y = -202.413 + 620.411 D_3$	7	.80	31.0 - 90.4 (cm)
Branches, 0.6 - 2.5 cm	ODW-kg	$Y = 13.632 + 2.784 D_3 CL_m$	7	.85	31.0 - 90.4 (cm)
Branches, < 0.6 & foliage	ODW-kg	$Y = -2.995 + 8.696 D_3 CL_m$	7	.95	31.0 - 90.4 (cm)
Tree, a.s. 30 cm	ODW-kg	$\log Y = 2.231 + 0.907 \log D_7$	7	.99	31.0 - 90.4 (cm)
Stem, a.s. 30 cm, wood	ODW-kg	$\log Y = 2.047 + 0.953 \log D_7$	7	.99	31.0 - 90.4 (cm)
Stem, a.s. 30 cm, bark	ODW-kg	$\log Y = 3.096 + 1.327 \log BA$	7	.95	31.0 - 90.4 (cm)
Branches, > 2.5 cm	ODW-kg	$\log Y = 2.665 + 2.493 \log D_3$	7	.92	31.0 - 90.4 (cm)
Branches, 0.6 - 2.5 cm	ODW-kg	$\log Y = 0.861 + 0.760 \log D_3 CL_m$	7	.95	31.0 - 90.4 (cm)
Branches, < 0.6 cm & foliage	ODW-kg	$\log Y = 0.879 + 1.038 \log D_3 CL_m$	7	.97	31.0 - 90.4 (cm)
Fir					
Amabilis (Pacific Silver) (Krumlik, 1974)					
Tree, a.s. 30.5 cm	ODW-kg	$Y = 176.567 + 128.139 D_7$	7	.99	15.24 - 76.20 (cm)
Bole, a.s. 30.5 cm t. 2.5 cm, wood	ODW-kg	$Y = 80.324 + 90.478 D_7$	7	.99	15.24 - 76.20 (cm)
Bole, a.s. 30.5 cm t. 2.5 cm, bark	ODW-kg	$Y = 17.753 + 18.763 D_7$	7	.96	15.24 - 76.20 (cm)
Branches, > 2.5 cm	ODW-kg	$Y = 1.041 + 11.625 D_7$	7	.80	15.24 - 76.20 (cm)
Tree, a.s. 30.5 cm	ODW-kg	$Y = -2029.050 + 6775.640 D_3$	7	.97	15.24 - 76.20 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, wood	ODW-kg	$Y = -1467.720 + 4769.210 D_3$	7	.97	15.24 - 76.20 (cm)
Branches, > 2.5 cm	ODW-kg	$Y = -202.413 + 620.411 D_3$	7	.80	15.24 - 76.20 (cm)
Branches, .64 - 2.5 cm	ODW-kg	$Y = -7.558 + 103.675 D_3$	7	.86	15.24 - 76.20 (cm)
Branches, < .64 cm & foliage	ODW-kg	$Y = -64.847 + 316.410 D_3$	7	.91	15.24 - 76.20 (cm)
Branches, .64 - 2.5 cm	ODW-kg	$Y = 13.632 + 2.784 D_3 CL_m$	7	.85	15.24 - 76.20 (cm)
Branches, < .64 cm & foliage	ODW-kg	$Y = -2.995 + 8.696 D_3 CL_m$	7	.95	15.24 - 76.20 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, wood	ODW-kg	$Y = -213.497 + 5089.520 BA$	7	.99	15.24 - 76.20 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, bark	ODW-kg	$Y = -42.324 + 1052.280 BA$	7	.95	15.24 - 76.20 (cm)
Branches, > 2.5 cm	ODW-kg	$Y = -39.770 + 663.778 BA$	7	.82	15.24 - 76.20 (cm)

Appendix

SPECIES (Reference)	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
Fir					
Amabilis (Pacific Silver) (Krumlik, 1974)					
..... cont'd					
Branches, < .64 cm & foliage	ODW-kg	$Y = 21.947 + 325.859 \text{ BA}$	7	.87	15.24 — 76.20 (cm)
Branches, > 2.5 cm	ODW-kg	$Y = 22.025 + 0.490 \text{ BA } H_3 \text{ CL}_m$	7	.76	15.24 — 76.20 (cm)
Branches < .64 cm & foliage	ODW-kg	$Y = 50.450 + 0.247 \text{ BA } H_3 \text{ CL}_m$	7	.85	15.24 — 76.20 (cm)
Branches, .64 — 2.5 cm	ODW-kg	$Y = 0.950 + 0.299 (\text{BA } H_3) + (\text{CL}_m \text{ CW}_m)$	7	.81	15.24 — 76.20 (cm)
Branches, < .64 cm & foliage	ODW-kg	$Y = -39.661 + 0.918 (\text{BA } H_3) + (\text{CL}_m \text{ CW}_m)$	7	.87	15.24 — 76.20 (cm)
Branches, .64 — 2.5 cm	ODW-kg	$Y = -24.609 + 3.472 \text{ CL}_m$	7	.80	15.24 — 76.20 (cm)
Branches, < .64 cm & foliage	ODW-kg	$Y = -118.107 + 10.651 \text{ CL}_m$	7	.86	15.24 — 76.20 (cm)
Branches, .64 — 2.5 cm	ODW-kg	$Y = -1.833 + 0.211 \text{ CS}$	7	.79	15.24 — 76.20 (cm)
Branches, .64 — 2.5 cm	ODW-kg	$Y = -28.944 + 2.660 \text{ H}_3$	7	.88	15.24 — 76.20 (cm)
Tree, a.s. 30.5 cm	ODW-kg	$\log Y = 2.279 + 0.899 \log D_7$	7	.99	15.24 — 76.20 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, wood	ODW-kg	$\log Y = 2.047 + 0.953 \log D_7$	7	.99	15.24 — 76.20 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, bark	ODW-kg	$\log Y = 1.346 + 0.967 \log D_7$	7	.96	15.24 — 76.20 (cm)
Branches, > 2.5 cm	ODW-kg	$\log Y = 1.169 + 0.890 \log D_7$	7	.89	15.24 — 76.20 (cm)
Branches, .64 — 2.5 cm	ODW-kg	$\log Y = 1.218 + 0.481 \log D_7$	7	.92	15.24 — 76.20 (cm)
Branches, < .64 cm & foliage	ODW-kg	$\log Y = 1.374 + 0.648 \log D_7$	7	.92	15.24 — 76.20 (cm)
Branches, > 2.5 cm	ODW-kg	$\log Y = 0.631 + 0.771 \log D_7 \text{ CW}_m$	7	.86	15.24 — 76.20 (cm)
Branches, .64 — 2.5 cm	ODW-kg	$\log Y = 0.918 + 0.421 \log D_7 \text{ CW}_m$	7	.91	15.24 — 76.20 (cm)
Branches, < .64 cm & foliage	ODW-kg	$\log Y = 0.983 + 0.561 \log D_7 \text{ CW}_m$	7	.89	15.24 — 76.20 (cm)
Tree, a.s. 30.5 cm	ODW-kg	$\log Y = 3.779 + 2.473 \log D_3$	7	.99	15.24 — 76.20 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, wood	ODW-kg	$\log Y = 3.636 + 2.618 \log D_3$	7	.99	15.24 — 76.20 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, bark	ODW-kg	$\log Y = 2.957 + 2.654 \log D_3$	7	.95	15.24 — 76.20 (cm)
Branches, > 2.5 cm	ODW-kg	$\log Y = 2.665 + 2.493 \log D_3$	7	.92	15.24 — 76.20 (cm)
Branches, .64 — 2.5 cm	ODW-kg	$\log Y = 2.019 + 1.317 \log D_3$	7	.91	15.24 — 76.20 (cm)
Branches, < .64 cm & foliage	ODW-kg	$\log Y = 2.457 + 1.789 \log D_3$	7	.92	15.24 — 76.20 (cm)
Branches, .64 — 2.5 cm	ODW-kg	$\log Y = 0.861 + 0.760 \log D_3 \text{ CL}_m$	7	.95	15.24 — 76.20 (cm)
Branches, < .64 cm & foliage	ODW-kg	$\log Y = 0.879 + 1.038 \log D_3 \text{ CL}_m$	7	.97	15.24 — 76.20 (cm)

Appendix

SPECIES (Reference) Equation Title	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Fir</u> Amabilis (Pacific Silver) (Krumlik, 1974) ... cont'd					
Branches, .64 - 2.5 cm	ODW-kg	$\log Y = 0.423 + 0.521 \log D_3 (CL_m)^2$	7	.95	15.24 - 76.20 (cm)
Branches, < .64 cm & foliage	ODW-kg	$\log Y = 0.276 + 0.713 \log D_3 (CL_m)^2$	7	.97	15.24 - 76.20 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, bark	ODW-kg	$\log Y = 3.096 + 1.327 \log 8A$	7	.95	15.24 - 76.20 (cm)
Branches, .64 - 2.5 cm	ODW-kg	$\log Y = -1.002 + 1.133 \log CS$	7	.90	15.24 - 76.20 (cm)
<u>Fir</u> Balsam (Baskerville, 1965) Tree	FW-lbs	$\log Y = 0.343 + 2.68 \log D_1$	98	.99 (r)	2.5 - 25.0 (cm)
Tree	ODW-lbs	$\log Y = 0.086 + 2.53 \log D_1$	101	.96 (r)	2.5 - 25.0 (cm)
Stem wood	ODW-lbs	$\log Y = 0.062 + 2.28 \log D_1$	101	.96 (r)	2.5 - 25.0 (cm)
Stem bark	ODW-lbs	$\log Y = -0.916 + 2.47 \log D_1$	101	.95 (r)	2.5 - 25.0 (cm)
Branches	ODW-lbs	$\log Y = -1.294 + 3.22 \log D_1$	101	.95 (r)	2.5 - 25.0 (cm)
Foliage	ODW-lbs	$\log Y = -1.258 + 3.21 \log D_1$	101	.98 (r)	2.5 - 25.0 (cm)
Cones (Dom. & Codom. only)	ODW-lbs	$\log_{100} Y = -0.625 + 3.20 \log D_1$	35	.70 (r)	10 - 25 (cm)
Dead branches	ODW-lbs	$\log_{10} Y = 0.226 + 2.11 \log D_1$	90	.80 (r)	2.5 - 25.0 (cm)
<u>Fir</u> Balsam (open grown) (Honer, 1971) Stem, s 6", wood	ODW-lbs	$\ln Y = -2.0029 + 1.6430 \ln D_1 + 0.8823 \ln H_1$	40	.99	2.5 - 40.0 (cm)
Stem, s 6", bark	ODW-lbs	$\ln Y = -1.3735 + 2.1140 \ln D_1$	40	.98	2.5 - 40.0 (cm)
Stem, s 6", wood and branch wood (1-3")	ODW-lbs	$\ln Y = -0.0501 + 2.2921 \ln D_1$	40	.99	2.5 - 40.0 (cm)
Foliage	ODW-lbs	$\ln Y = 0.4408 + 1.7853 \ln D_1$	40	.97	2.5 - 40.0 (cm)
Tree, s 6"	ODW-lbs	$\ln Y = 1.3479 + 2.0543 \ln D_1$	40	.99	2.5 - 40.0 (cm)
Roots and stump s 6"	ODW-lbs	$\ln Y = 0.0035 + 2.0027 \ln D_1$	40	.93	2.5 - 40.0 (cm)
Tree tot.	ODW-lbs	$\ln Y = 1.6067 + 2.0344 \ln D_1$	40	.98	2.5 - 40.0 (cm)
Branch wood, bark, twigs	ODW-lbs	$\ln Y = 0.0707 + 2.0964 \ln D_1$	40	.98	2.5 - 40.0 (cm)

Appendix

SPECIES (Reference) Equation Title	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Fir</u>					
Balsam (forest grown) (Honer, 1971)					
Stem, s 6", wood	ODW-lbs	$\ln Y = -4.3020 + 1.6350 \ln D_1 + 1.4934 \ln H_1$	40	.99	2.5 - 40.0 (cm)
Stem, s 6", bark	ODW-lbs	$\ln Y = -5.7100 + 1.8333 \ln D_1 + 1.3178 \ln H_1$	40	.94	2.5 - 40.0 (cm)
Stem, s 6", wood and branch wood (1-3")	ODW-lbs	$\ln Y = -4.1291 + 1.6997 \ln D_1 + 1.4210 \ln H_1$	40	.99	2.5 - 40.0 (cm)
Foliage	ODW-lbs	$\ln Y = 5.4899 + 3.5856 \ln D_1 - 2.3816 \ln H_1$	40	.69	2.5 - 40.0 (cm)
Tree, s 6"	ODW-lbs	$\ln Y = 0.4441 + 2.4975 \ln D_1$	40	.97	2.5 - 40.0 (cm)
Roots and stump, s 3"	ODW-lbs	$\ln Y = -1.0678 + 2.4613 \ln D_1$	40	.90	2.5 - 40.0 (cm)
Tree tot.	ODW-lbs	$\ln Y = 0.6538 + 2.4872 \ln D_1$	40	.97	2.5 - 40.0 (cm)
Branch wood, bark and twigs	ODW-lbs	$\ln Y = 5.6263 + 4.4912 \ln D_1 - 2.7786 \ln H_1$	40	.78	2.5 - 40.0 (cm)
<u>Fir</u>					
Balsam (Keen, 1963)					
Bole, s*, t 3", wood	FW-lbs	$\log Y = 0.108 + 2.673 \log D_1$	64	.95 (r)	10.0 - 40.0 (cm)
Bole, s*, t 3"	FW-lbs	$\log Y = 0.212 + 2.650 \log D_1$	201	.96 (r)	10.0 - 40.0 (cm)
Tree, s*	FW-lbs	$\log Y = 0.639 + 2.382 \log D_1$	138	.97 (r)	10.0 - 40.0 (cm)
Bole, s*, t 3"	FW-lbs	$\log Y = -0.7809 + 2.021 \log D_1 + 0.919 \log H_1$	192	.98 (r)	10.0 - 40.0 (cm)
Bole, s*, t 3" (* = D ₊)	FW-lbs	$\log Y = -0.7494 + 1.983 \log D_1 + 0.879 \log H_1$	64	.98 (r)	10.0 - 40.0 (cm)
Slash	FW-lbs	$\log Y = 0.610 + 1.872 \log D_1$	138	.86 (r)	10.0 - 40.0 (cm)
<u>Fir</u>					
Balsam (Young et al., 1964)					
Tree tot.	FW-lbs	$\ln Y = .5882 + 2.2234 \ln D_1 + .3390 \ln H_1$	23	.97	15.5 - 33.5 (cm)
Tree tot.	ODW-lbs	22% of FW from above equation	23	.97	15.5 - 33.5 (cm)
Tree tot. - Roots < 4"	FW-lbs	$\ln Y = -.0260 + 2.0530 \ln D_1 + .5656 \ln H_1$	23	.97	15.5 - 33.5 (cm)
Tree tot. - Roots < 4"	ODW-lbs	23% of FW from above equation	23	.97	15.5 - 33.5 (cm)
Tree a.s. 6"	FW-lbs	$\ln Y = .0511 + 2.1679 \ln D_1 + .4292 \ln H_1$	23	.96	15.5 - 33.5 (cm)
Tree a.s. 6"	ODW-lbs	21% of FW from above equation	23	.96	15.5 - 33.5 (cm)

Appendix

SPECIES (Reference)	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST-HEIGHT IN CM
<u>Fir</u>					
Balsam (Young et al., 1964)					
.... cont'd					
Stem a.s. 6"	FW-lbs	$\ln Y = 1.5589 + 1.8801 \ln D_1 + .9332 \ln H_1$	23	.98	15.5 — 33.5 (cm)
Stem a.s. 6"	ODW-lbs	33% of FW from above equation	23	.98	15.5 — 33.5 (cm)
Bole a.s. 6", t. 4"	FW-lbs	$\ln Y = -2.0858 + 2.0215 \ln D_1 + .9670 \ln H_1$	23	.98	15.5 — 33.5 (cm)
Bole a.s. 6", t. 4"	ODW-lbs	33% of FW from above equation	23	.98	15.5 — 33.5 (cm)
All branches	FW-lbs	$\ln Y = 2.2870 + 3.2160 \ln D_1 - 1.0895 \ln H_1$	23	.85	15.5 — 33.5 (cm)
All branches	ODW-lbs	8% of FW from above equation	23	.85	15.5 — 33.5 (cm)
Branches < 1"	FW-lbs	$\ln Y = 1.9971 + 2.7950 \ln D_1 - .8048 \ln H_1$	23	.82	15.5 — 33.5 (cm)
Branches < 1"	ODW-lbs	7% of FW from above equation	23	.82	15.5 — 33.5 (cm)
Stump 6" and roots > 1"	FW-lbs	$\ln Y = .3669 + 2.6447 \ln D_1 - .2645 \ln H_1$	23	.92	15.5 — 33.5 (cm)
Stump 6" and roots > 1"	ODW-lbs	30% of FW from above equation	23	.92	15.5 — 33.5 (cm)
Roots > 1"	FW-lbs	$\ln Y = .8681 + 2.7785 \ln D_1 - .5456 \ln H_1$	23	.92	15.5 — 33.5 (cm)
Roots > 1"	ODW-lbs	28% of FW from above equation	23	.92	15.5 — 33.5 (cm)
Roots < 1"	FW-lbs	$\ln Y = -1.1186 + 1.6306 \ln D_1 + .3564 \ln H_1$	23	.50	15.5 — 33.5 (cm)
Roots < 1"	ODW-lbs	12% of FW from above equation	23	.50	15.5 — 33.5 (cm)
<u>Fir</u>					
Balsam (Young and Carpenter, 1967)					
Foliage	FW-g	$\ln Y = 3.141 + 1.963 \ln H_1$	14	.95	0.0 — 75 (mm)
Branches	FW-g	$\ln Y = 2.476 + 1.948 \ln H_1$	14	.93	0.0 — 75 (mm)
Stem	FW-g	$\ln Y = 1.095 + 2.871 \ln H_1$	14	.95	0.0 — 75 (mm)
Roots	FW-g	$\ln Y = 2.332 + 2.165 \ln H_1$	14	.94	0.0 — 75 (mm)
Tree tot.	FW-g	$\ln Y = 3.909 + 2.160 \ln H_1$	14	.97	0.0 — 75 (mm)
Foliage	ODW-g	$\ln Y = 2.324 + 1.941 \ln H_1$	14	.95	0.0 — 75 (mm)
Branches	ODW-g	$\ln Y = 1.851 + 1.934 \ln H_1$	14	.93	0.0 — 75 (mm)
Stem	ODW-g	$\ln Y = 2.273 + 2.893 \ln H_1$	14	.94	0.0 — 75 (mm)
Roots	ODW-g	$\ln Y = 1.569 + 2.122 \ln H_1$	14	.93	0.0 — 75 (mm)
Tree tot.	ODW-g	$\ln Y = 3.147 + 2.145 \ln H_1$	14	.96	0.0 — 75 (mm)

Appendix

SPECIES (Reference) Equation Title	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Fir</u>					
Douglas (Kurucz, 1969) Tree	ODW-lbs	$Y = 0.059265 D_6$	112	.96	2.5 - 20.0 (cm)
Stem - wood	ODW-lbs	$Y = 0.044119 D_6$	112	.98	2.5 - 20.0 (cm)
Stem - bark	ODW-lbs	$Y = 0.007908 D_6$	112	.98	2.5 - 20.0 (cm)
Crown	ODW-lbs	$Y = 0.080017 DC_3$	112	.59	2.5 - 20.0 (cm)
Branches > 3/4 in	ODW-lbs	$Y = 0.048776 DC_3$	112	.58	2.5 - 20.0 (cm)
Branches 1/4 - 3/4 in	ODW-lbs	$Y = 0.006931 DC_3$	112	.61	2.5 - 20.0 (cm)
Branches < 1/4 in	ODW-lbs	$Y = 0.007999 DC_3$	112	.58	2.5 - 20.0 (cm)
Foliage	ODW-lbs	$Y = 0.016310 DC_3$	112	.56	2.5 - 20.0 (cm)
<u>Fir</u>					
Subalpine (Kimmins, 1974) Bole, a.s. 30.5 cm, t. 2.5 cm, wood	ODW-kg	$\log Y = 1.467 + 1.63 \log D_3 + 1.299 \log H_3$	8	.99	14.1 - 60.0 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, bark	ODW-kg	$\log Y = 0.454 + 1.336 \log D_3 + 1.476 \log H_3$	8	.99	14.1 - 60.0 (cm)
Branches > 2.5 cm	ODW-kg	$\log Y = 9.960 + 7.882 \log D_3 - 3.713 \log H_3$	8	.83	14.1 - 60.0 (cm)
Branches, .64 - 2.5 cm	ODW-kg	$\log Y = 1.555 - 1.216 \log D_3 CL_m + 0.104 \log D_3 CL_m$	8	.66	14.1 - 60.0 (cm)
Branches, < .64 cm & foliage	ODW-kg	$\log Y = 3.767 + 1.893 \log D_3 - 0.859 \log H_3$	8	.89	14.1 - 60.0 (cm)
<u>Hemlock</u> sp. (Young and Carpenter, 1967)					
Foliage	FW-g	$\ln Y = 2.197 + 2.147 \ln H_1$	9	.98	0.0 - 75 (mm)
Branches	FW-g	$\ln Y = 1.374 + 2.306 \ln H_1$	9	.97	0.0 - 75 (mm)
Stem	FW-g	$\ln Y = -1.345 + 3.605 \ln H_1$	9	.94	0.0 - 75 (mm)
Roots	FW-g	$\ln Y = 1.229 + 2.448 \ln H_1$	9	.98	0.0 - 75 (mm)
Tree tot.	FW-g	$\ln Y = 2.789 + 2.476 \ln H_1$	9	.98	0.0 - 75 (mm)
Foliage	ODW-g	$\ln Y = 1.266 + 2.192 \ln H_1$	9	.98	0.0 - 75 (mm)
Branches	ODW-g	$\ln Y = 0.779 + 2.320 \ln H_1$	9	.98	0.0 - 75 (mm)

Appendix

SPECIES (Reference)	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Hemlock</u> sp. (Young and Carpenter, 1967) cont'd					
Stem	ODW-g	$\ln Y = -1.414 + 3.343 \ln H_1$	9	.94	0.0 - 75 (mm)
Roots	ODW-g	$\ln Y = 0.549 + 2.405 \ln H_1$	9	.98	0.0 - 75 (mm)
Tree tot.	ODW-g	$\ln Y = 2.035 + 2.476 \ln H_1$	9	.98	0.0 - 75 (mm)
<u>Hemlock</u> Eastern (Young et al., 1964)					
Tree tot.	FW-lbs	$\ln Y = .6277 + 2.2234 \ln D_1 + .3390 \ln H_1$	28	.98	14.0 - 37.8 (cm)
Tree tot.	ODW-lbs	31% of FW from above equation	28	.98	14.0 - 37.8 (cm)
Tree tot. - Roots < 4"	FW-lbs	$\ln Y = -.3640 + 2.0634 \ln D_1 + .6564 \ln H_1$	28	.98	14.0 - 37.8 (cm)
Tree tot. - Roots < 4"	ODW-lbs	32% of FW from above equation	28	.98	14.0 - 37.8 (cm)
Tree a.s. 6"	FW-lbs	$\ln Y = .1798 + 2.1679 \ln D_1 + .4292 \ln H_1$	28	.98	14.0 - 37.8 (cm)
Tree a.s. 6"	ODW-lbs	31% of FW from above equation	28	.98	14.0 - 37.8 (cm)
Stem a.s. 6"	FW-lbs	$\ln Y = -1.5128 + 1.8801 \ln D_1 + .9332 \ln H_1$	28	.83	14.0 - 37.8 (cm)
Stem a.s. 6"	ODW-lbs	38% of FW from above equation	28	.83	14.0 - 37.8 (cm)
Bole a.s. 6", t. 4"	FW-lbs	$\ln Y = -2.0319 + 2.0215 \ln D_1 + .9670 \ln H_1$	28	.98	14.0 - 37.8 (cm)
Bole a.s. 6", t. 4"	ODW-lbs	38% of FW from above equation	28	.98	14.0 - 37.8 (cm)
All branches	FW-lbs	$\ln Y = 2.1610 + 2.8863 \ln D_1 - .8043 \ln H_1$	28	.89	14.0 - 37.8 (cm)
All branches	ODW-lbs	14% of FW from above equation	28	.89	14.0 - 37.8 (cm)
Branches < 1 in	FW-lbs	$\ln Y = 1.7234 + 2.3289 \ln D_1 - .4612 \ln H_1$	28	.80	14.0 - 37.8 (cm)
Branches < 1 in	ODW-lbs	10% of FW from above equation	28	.80	14.0 - 37.8 (cm)
Stump 6" and roots > 1 in	FW-lbs	$\ln Y = -2.7994 + 2.6523 \ln D_1 + .4997 \ln H_1$	28	.97	14.0 - 37.8 (cm)
Stump 6" and roots > 1 in	ODW-lbs	39% of FW from above equation	28	.97	14.0 - 37.8 (cm)
Roots > 1 in	FW-lbs	$\ln Y = -3.9057 + 2.6756 \ln D_1 + .6633 \ln H_1$	28	.95	14.0 - 37.8 (cm)
Roots > 1 in	ODW-lbs	40% of FW from above equation	28	.95	14.0 - 37.8 (cm)
Roots < 1 in	FW-lbs	$\ln Y = -1.4966 + 1.6306 \ln D_1 + .3564 \ln H_1$	28	.59	14.0 - 37.8 (cm)
Roots < 1 in	ODW-lbs	11% of FW from above equation	28	.59	14.0 - 37.8 (cm)

Appendix

SPECIES (Reference) Equation Title	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND-LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Hemlock</u> Mountain (Krumlik and Kimmins, 1973)					
Tree, a.s. 30 cm	ODW-kg	$Y = 386.715 + 117.631 D_7$	5	.99	43.7 - 76.2 (cm)
Stem, a.s. 30 cm, wood	ODW-kg	$Y = 266.160 + 86.292 D_7$	5	.99	43.7 - 76.2 (cm)
Stem, a.s. 30 cm, bark	ODW-kg	$Y = -15.193 + 1282.740 BA$	5	.99	43.7 - 76.2 (cm)
Branches > 2.5 cm	ODW-kg	$Y = 22.048 + 1.378 D_7 CW_m$	5	.99	43.7 - 76.2 (cm)
Branches < .6 cm & foliage	ODW-kg	$Y = -20.637 + 9.679 D_3 CL_m$	5	.97	43.7 - 76.2 (cm)
Branches < .6 cm	ODW-kg	$Y = -18.780 + 4.700 D_3 CL_m$	5	.99	43.7 - 76.2 (cm)
Foliage	ODW-kg	$Y = -1.857 + 4.980 D_3 CL_m$	5	.88	43.7 - 76.2 (cm)
Tree, a.s. 30 cm	ODW-kg	$\log Y = 2.464 + .742 \log D_7$	5	.99	43.7 - 76.2 (cm)
Stem, a.s. 30 cm, wood	ODW-kg	$\log Y = 2.319 + .746 \log D_7$	5	.98	43.7 - 76.2 (cm)
Stem, a.s. 30 cm, bark	ODW-kg	$\log Y = 3.109 + 1.039 \log BA$	5	.98	43.7 - 76.2 (cm)
Branches > 2.5 cm	ODW-kg	$\log Y = 2.800 + 3.074 \log D_3$	5	.99	43.7 - 76.2 (cm)
Branches < .6 cm & foliage	ODW-kg	$\log Y = 0.567 + 1.297 \log D_3 CL_m$	5	.96	43.7 - 76.2 (cm)
Branches < .6 cm	ODW-kg	$\log Y = -1.272 + 1.163 \log D_3 (CL_m)^2$	5	.99	43.7 - 76.2 (cm)
Foliage	ODW-kg	$\log Y = -0.537 + 0.936 \log D_3 (CL_m)^2$	5	.90	43.7 - 76.2 (cm)
<u>Hemlock</u> Mountain (Krumlik, 1974)					
Tree, a.s. 30.6 cm	ODW-kg	$Y = 366.484 + 131.667 D_7$	5	.99	15.24 - 76.20 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, wood	ODW-kg	$Y = 266.160 + 86.292 D_7$	5	.99	15.24 - 76.20 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, bark	ODW-kg	$Y = 58.359 + 24.286 D_7$	5	.97	15.24 - 76.20 (cm)
Branches, > 2.5 cm	ODW-kg	$Y = -32.176 + 14.498 D_7$	5	.94	15.24 - 76.20 (cm)
Branches, < .64 cm & foliage	ODW-kg	$Y = 50.185 + 5.564 D_7$	5	.87	15.24 - 76.20 (cm)
Branches, < .64 cm	ODW-kg	$Y = 15.286 + 2.727 D_7$	5	.91	15.24 - 76.20 (cm)
Foliage	ODW-kg	$Y = 34.899 + 2.837 D_7$	5	.78	15.24 - 76.20 (cm)
Tree, a.s. 30.5 cm	ODW-kg	$Y = -1904.550 + 6526.130 D_3$	5	.98	15.24 - 76.20 (cm)

Appendix

SPECIES	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Hemlock</u>					
<u>Mountain</u>					
(Krumlik, 1974)					
.... cont'd					
Bole, a.s. 30.5 cm, t. 2.5 cm, wood	ODW-kg	$Y = -1194.950 + 4231.780 D_3$	5	.95	15.24 - 76.20 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, bark	ODW-kg	$Y = -369.519 + 1218.670 D_3$	5	.98	15.24 - 76.20 (cm)
Branches, > 2.5 cm	ODW-kg	$Y = -290.516 + 732.353 D_3$	5	.96	15.24 - 76.20 (cm)
Branches, < .64 cm & foliage	ODW-kg	$Y = -51.448 + 285.195 D_3$	5	.92	15.24 - 76.20 (cm)
Branches, < .64 cm	ODW-kg	$Y = -34.157 + 139.157 D_3$	5	.95	15.24 - 76.20 (cm)
Foliage	ODW-kg	$Y = -17.296 + 146.040 D_3$	5	.83	15.24 - 76.20 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, wood	ODW-kg	$Y = 42.538 + 4430.220 BA$	5	.95	15.24 - 76.20 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, bark	ODW-kg	$Y = -15.193 + 1282.740 BA$	5	.99	15.24 - 76.20 (cm)
Branches, > 2.5 cm	ODW-kg	$Y = -79.402 + 776.993 BA$	5	.98	15.24 - 76.20 (cm)
Branches, < .64 cm & foliage	ODW-kg	$Y = 33.897 + 291.988 BA$	5	.88	15.24 - 76.20 (cm)
Branches, > 2.5 cm	ODW-kg	$Y = 22.048 + 1.378 D_7 CW_m$	5	.99	15.24 - 76.20 (cm)
Branches, > 2.5 cm	ODW-kg	$Y = -.0586 + 0.504 CV$	5	.97	15.24 - 76.20 (cm)
Branches, < .64 cm & foliage	ODW-kg	$Y = -20.637 + 9.679 D_3 CL_m$	5	.97	15.24 - 76.20 (cm)
Branches, < .64 cm	ODW-kg	$Y = -18.780 + 4.700 D_3 CL_m$	5	.99	15.24 - 76.20 (cm)
Foliage	ODW-kg	$Y = -1.857 + 4.980 D_3 CL_m$	5	.88	15.24 - 76.20 (cm)
Branches, < .64 cm & foliage	ODW-kg	$Y = 53.070 + 0.277 BA H_3 CL_m$	5	.90	15.24 - 76.20 (cm)
Branches, < .64 cm	ODW-kg	$Y = 16.662 + 0.136 BA H_3 CL_m$	5	.94	15.24 - 76.20 (cm)
Foliage	ODW-kg	$Y = 36.408 + 0.141 BA H_3 CL_m$	5	.81	15.24 - 76.20 (cm)
Branches, < .64 cm & foliage	ODW-kg	$Y = 22.995 + 0.574 (BA H_3) + (CL_m CW_m)$	5	.84	15.24 - 76.20 (cm)
Branches, < .64 cm	ODW-kg	$Y = 0.143 + 0.292 (BA H_3) + (CL_m CW_m)$	5	.94	15.24 - 76.20 (cm)
Branches, < .64 cm	ODW-kg	$Y = 0.027 + 0.196 CS$	5	.93	15.24 - 76.20 (cm)
Tree, a.s. 30.5 cm	ODW-kg	$\log Y = 2.469 + 0.769 \log D_7$	5	.99	15.24 - 76.20 (cm)
Bole, 30.5 cm, t. 2.5 cm, bark	ODW-kg	$\log Y = 1.701 + 0.778 \log D_7$	5	.98	15.24 - 76.20 (cm)

Appendix

SPECIES (Reference)	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Hemlock</u>					
<u>Mountain</u>					
(Krumlik, 1974)					
..... cont'd					
Bole, a.s. 30.5 cm. t. 2.5 cm, wood	ODW-kg	$\log Y = 2.319 + 0.746 \log D_7$	5	.98	15.24 — 76.20 (cm)
Branches, > 2.5 cm	ODW-kg	$\log Y = 0.894 + 1.151 \log D_7$	5	.96	15.24 — 76.20 (cm)
Branches, < .64 cm	ODW-kg	$\log Y = 0.909 + 0.724 \log D_7$	5	.94	15.24 — 76.20 (cm)
Branches, < .64 cm & foliage	ODW-kg	$\log Y = 0.885 + 1.558 \log D_7 - 0.036 \log D_7$	5	.99	15.24 — 76.20 (cm)
Foliage	ODW-kg	$\log Y = 0.598 + 1.7174 \log D_7 + 0.045 \log D_7$	5	.98	15.24 — 76.20 (cm)
Tree, a.s. 30.5 cm	ODW-kg	$\log Y = 3.733 + 2.016 \log D_3$	5	.99	15.24 — 76.20 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, wood	ODW-kg	$\log Y = 3.544 + 1.947 \log D_3$	5	.97	15.24 — 76.20 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, bark	ODW-kg	$\log Y = 3.000 + 2.078 \log D_3$	5	.98	15.24 — 76.20 (cm)
Branches, > 2.5 cm	ODW-kg	$\log Y = 2.800 + 3.074 \log D_3$	5	.99	15.24 — 76.20 (cm)
Branches, < .64 cm & foliage	ODW-kg	$\log Y = 2.434 + 1.651 \log D_3$	5	.88	15.24 — 76.20 (cm)
Branches, < .64 cm	ODW-kg	$\log Y = 2.099 + 1.895 \log D_3$	5	.93	15.24 — 76.20 (cm)
Foliage	ODW-kg	$\log Y = 2.168 + 1.486 \log D_3$	5	.80	15.24 — 76.20 (cm)
Bole, a.s. 30.5 cm, t.2.5 cm, bark	ODW-kg	$\log Y = 3.109 + 1.039 \log BA$	5	.98	15.24 — 76.20 (cm)
Branches, > 2.5 cm	ODW-kg	$\log Y = 1.291 + 0.005 \log (BA H_3) + (CL_m CW_m)$	5	.95	15.24 — 76.20 (cm)
Branches, < .64 cm & foliage	ODW-kg	$\log Y = 0.567 + 1.297 \log D_3 CL_m$	5	.96	15.24 — 76.20 (cm)
Branches, < .64 cm	ODW-kg	$\log Y = -2.780 + 1.474 \log D_3 CL_m$	5	.99	15.24 — 76.20 (cm)
Foliage	ODW-kg	$\log Y = 0.478 + 1.176 \log D_3 CL_m$	5	.99	15.24 — 76.20 (cm)
Branches, < .64 cm & foliage	ODW-kg	$\log Y = -0.542 + 1.029 \log D_3 (CL_m)^2$	5	.97	15.24 — 76.20 (cm)
Branches, < .64 cm	ODW-kg	$\log Y = -1.272 + 1.163 \log D_3 (CL_m)^2$	5	.99	15.24 — 76.20 (cm)
Foliage	ODW-kg	$\log Y = -0.537 + 0.936 \log D_3 (CL_m)^2$	5	.90	15.24 — 76.20 (cm)
Branches, > 2.5 cm	ODW-kg	$\log Y = 0.647 + 0.785 \log D_7 CW_m$	5	.99	15.24 — 76.20 (cm)
Branches, < .64 cm & foliage	ODW-kg	$\log Y = 1.290 + 0.415 \log D_7 CW_m$	5	.86	15.24 — 76.20 (cm)
Branches, < .64 cm	ODW-kg	$\log Y = 0.122 + 0.971 \log D_7 CW_m - 0.003 \log D_7 CW_m$	5	.99	15.24 — 76.20 (cm)
Branches, > 2.5 cm	ODW-kg	$\log Y = 1.302 + 0.003 \log CS$	5	.95	15.24 — 76.20 (cm)

Appendix

SPECIES (Reference)	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Hemlock</u>					
<u>Western</u>					
(Krumhik, 1974)					
Tree, a.s. 30.5 cm	ODW-kg	$Y = 79.458 + 136.626 D_7$	8	.96	15.24 — 127.0 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, wood	ODW-kg	$Y = 48.856 + 90.959 D_7$	8	.96	15.24 — 127.0 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, wood	ODW-kg	$Y = -9.236 + 3342.380 BA$	8	.98	15.24 — 127.0 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, bark	ODW-kg	$Y = -10.054 + 879.140 BA$	8	.94	15.24 — 127.0 (cm)
Foliage	ODW-kg	$Y = -1.002 + 279.218 BA$	8	.86	15.24 — 127.0 (cm)
Branches, > 2.5 cm	ODW-kg	$Y = 2.919 + 1.042 BA H_3 CL_m$	8	.88	15.24 — 127.0 (cm)
Branches, .64 — 2.5 cm	ODW-kg	$Y = 18.807 + 13.101 BA H_3$	8	.71	15.24 — 127.0 (cm)
Branches, < .64 cm & foliage	ODW-kg	$Y = 7.680 + 14.039 BA H_3$	8	.85	15.24 — 127.0 (cm)
Foliage	ODW-kg	$Y = 3.595 + 9.803 BA H_3$	8	.87	15.24 — 127.0 (cm)
Branches, < .64 cm	ODW-kg	$Y = -0.618 + 0.160 (BA H_3) + (CL_m CW_m)$	8	.85	15.24 — 127.0 (cm)
Foliage	ODW-kg	$Y = -3.515 + 0.325 (BA H_3) + (CL_m CW_m)$	8	.77	15.24 — 127.0 (cm)
Branches, > 2.5 cm	ODW-kg	$Y = -27.147 + 13.161 D_3 CL_m$	8	.92	15.24 — 127.0 (cm)
Branches, .64 — 2.4 cm	ODW-kg	$Y = 8.504 + 7.119 D_3 CL_m$	8	.76	15.24 — 127.0 (cm)
Branches, < .64 cm & foliage	ODW-kg	$Y = -2.601 + 7.480 D_3 CL_m$	8	.88	15.24 — 127.0 (cm)
Branches, < .64 cm	ODW-kg	$Y = 0.772 + 2.296 D_3 CL_m$	8	.79	15.24 — 127.0 (cm)
Foliage	ODW-kg	$Y = -3.340 + 5.176 D_3 CL_m$	8	.88	15.24 — 127.0 (cm)
Branches, > 2.5 cm	ODW-kg	$Y = 1.759 + 2.554 D_7 CW_m$	8	.88	15.24 — 127.0 (cm)
Branches, .64 — 2.5 cm	ODW-kg	$Y = -18.780 + 3.565 D_7 + CL_m$	8	.73	15.24 — 127.0 (cm)
Branches, > 2.5 cm	ODW-kg	$Y = -8.717 + 0.398 CV$	8	.89	15.24 — 127.0 (cm)
Branches, < .64 cm & foliage	ODW-kg	$Y = 8.385 + 0.222 CV$	8	.82	15.24 — 127.0 (cm)
Branches, < .64 cm	ODW-kg	$Y = 3.468 + 0.074 CV$	8	.86	15.24 — 127.0 (cm)
Branches, > 2.5 cm	ODW-kg	$Y = -30.646 + 0.544 CS$	8	.87	15.24 — 127.0 (cm)
Branches, .64 — 2.5 cm	ODW-kg	$Y = 6.726 + 0.293 CS$	8	.72	15.24 — 127.0 (cm)
Branches, < .64 cm & foliage	ODW-kg	$Y = -4.197 + 0.306 CS$	8	.82	15.24 — 127.0 (cm)

Appendix

SPECIES (Reference)	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Hemlock</u>					
<u>Western</u>					
(Krumlik, 1974)					
.... cont'd					
Branches, < .64 cm	ODW-kg	$Y = -0.633 + 0.101 CS$	8	.85	15.24 - 127.0 (cm)
Foliage	ODW-kg	$Y = -3.530 + 0.205 CS$	8	.77	15.24 - 127.0 (cm)
Branches, > 2.5 cm	ODW-kg	$Y = -99.310 + 9.100 CL_m$	8	.81	15.24 - 127.0 (cm)
Branches, < .64 cm & foliage	ODW-kg	$Y = -3.758 + 1.459 CW_m^2$	8	.86	15.24 - 127.0 (cm)
Branches, < .64 cm	ODW-kg	$Y = -0.621 + 0.486 CW_m^2$	8	.91	15.24 - 127.0 (cm)
Branches, .64 - 2.5 cm	ODW-kg	$Y = -72.493 + 5.647 H_3$	8	.87	15.24 - 127.0 (cm)
Branches, < .64 cm & foliage	ODW-kg	$Y = -82.759 + 5.695 H_3$	8	.93	15.24 - 127.0 (cm)
Branches, < .64 cm	ODW-kg	$Y = -23.751 + 1.74 H_3$	8	.83	15.24 - 127.0 (cm)
Foliage	ODW-kg	$Y = -58.907 + 3.946 H_3$	8	.94	15.24 - 127.0 (cm)
Tree, a.s. 30.5 cm	ODW-kg	$\log Y = 2.304 + 0.845 \log D_7$	8	.99	15.24 - 127.0 (cm)
Branches, > 2.5 cm	ODW-kg	$\log Y = 0.825 + 1.570 \log D_7$	8	.77	15.24 - 127.0 (cm)
Branches, .64 - 2.5 cm	ODW-kg	$\log Y = 1.427 + 0.620 \log D_7$	8	.90	15.24 - 127.0 (cm)
Branches, < .64 cm & foliage	ODW-kg	$\log Y = 0.125 + 0.774 \log D_7$	8	.92	15.24 - 127.0 (cm)
Foliage	ODW-kg	$\log Y = 0.951 + 1.022 \log D_7$	8	.92	15.24 - 127.0 (cm)
Branches, > 2.5 cm	ODW-kg	$\log Y = 0.072 + 1.231 \log D_7 CW_m$	8	.75	15.24 - 127.0 (cm)
Branches, < .64 cm & foliage	ODW-kg	$\log Y = 0.863 + 0.626 \log D_7 CW_m$	8	.96	15.24 - 127.0 (cm)
Branches, < .64 cm	ODW-kg	$\log Y = 0.560 + 0.451 \log D_7 CW_m$	8	.74	15.24 - 127.0 (cm)
Foliage	ODW-kg	$\log Y = 0.455 + 0.807 \log D_7 CW_m$	8	.92	15.24 - 127.0 (cm)
Tree, a.s. 30.5 cm	ODW-kg	$\log Y = 3.608 + 2.058 \log D_3$	8	.97	15.24 - 127.0 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, wood	ODW-kg	$\log Y = 3.455 + 2.120 \log D_3$	8	.98	15.24 - 127.0 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, bark	ODW-kg	$\log Y = 2.934 + 2.394 \log D_3$	8	.94	15.24 - 127.0 (cm)
Branches, > 2.5 cm	ODW-kg	$\log Y = 3.270 + 3.868 \log D_3$	8	.78	15.24 - 127.0 (cm)
Branches, .64 - 2.5 cm	ODW-kg	$\log Y = 2.366 + 1.477 \log D_3$	8	.85	15.24 - 127.0 (cm)
Branches, < .64 cm & foliage	ODW-kg	$\log Y = 2.420 + 1.837 \log D_3$	8	.86	15.24 - 127.0 (cm)

Appendix

SPECIES (Reference)	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Hemlock</u>					
<u>Western</u>					
(Krumlik, 1974)					
... cont'd					
Foliage	ODW-kg	$\log Y = 2.508 + 2.454 \log D_3$	8	.89	15.24 - 127.0 (cm)
Branches, .64 - 2.5 cm	ODW-kg	$\log Y = 0.936 + 1.007 \log D_3 CL_m$	8	.91	15.24 - 127.0 (cm)
Branches, < .64 cm & foliage	ODW-kg	$\log Y = 0.628 + 1.272 \log D_3 CL_m$	8	.96	15.24 - 127.0 (cm)
Branches, < .64 cm	ODW-kg	$\log Y = 0.422 + 0.931 \log D_3 CL_m$	8	.77	15.24 - 127.0 (cm)
Foliage	ODW-kg	$\log Y = -0.150 + 3.272 \log D_3 CL_m - 0.144 \log D_3 CL_m$	8	.97	15.24 - 127.0 (cm)
Branches, .64 - 2.5 cm	ODW-kg	$\log Y = 0.317 + 0.697 \log D_3 (CL_m)^2$	8	.86	15.24 - 127.0 (cm)
Branches, < .64 cm & foliage	ODW-kg	$\log Y = -0.165 + 0.888 \log D_3 (CL_m)^2$	8	.92	15.24 - 127.0 (cm)
Branches, < .64 cm	ODW-kg	$\log Y = -0.213 + 0.679 \log D_3 (CL_m)^2$	8	.80	15.24 - 127.0 (cm)
Foliage	ODW-kg	$\log Y = -0.816 + 1.114 \log D_3 (CL_m)^2$	8	.84	15.24 - 127.0 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, wood	ODW-kg	$\log Y = 2.112 + 0.370 \log D_7$	8	.99	15.24 - 127.0 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, bark	ODW-kg	$\log Y = 1.415 + 0.996 \log D_7$	8	.97	15.24 - 127.0 (cm)
Branches, .64 - 2.5 cm	ODW-kg	$\log Y = 1.177 + 0.499 \log D_7 CW_m$	8	.93	15.24 - 127.0 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, bark	ODW-kg	$\log Y = 3.060 + 1.197 \log BA$	8	.94	15.24 - 127.0 (cm)
Branches, < .64 cm	ODW-kg	$\log Y = 0.564 + 0.006 \log (BA H_3) + (CL_m CW_m)$	8	.83	15.24 - 127.0 (cm)
Branches, .64 - 2.5 cm	ODW-kg	$\log Y = 0.406 + 0.602 \log CV$	8	.69	15.24 - 127.0 (cm)
Branches, < .64 cm & foliage	ODW-kg	$\log Y = -0.053 + 0.767 \log CV$	8	.74	15.24 - 127.0 (cm)
Branches, < .64 cm	ODW-kg	$\log Y = 0.563 + 0.003 \log CS$	8	.83	15.24 - 127.0 (cm)
<u>Hemlock</u>					
<u>Western</u>					
(Krumlik and Kimmins, 1973)					
Tree, a.s. 30 cm	ODW-kg	$Y = 79.458 + 136.626 D_7$	8	.96	16.0 - 49.0 (cm)
Stem, a.s. 30 cm, wood	ODW-kg	$Y = 48.856 + 90.959 D_7$	8	.96	16.0 - 49.0 (cm)
Stem a.s. 30 cm, bark	ODW-kg	$Y = -10.054 + 879.140 BA$	8	.94	16.0 - 49.0 (cm)
Branches, > 2.5 cm	ODW-kg	$Y = 1.759 + 2.554 D_7 CW_m$	8	.88	16.0 - 49.0 (cm)

Appendix

SPECIES (Reference)	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Hemlock</u>					
<u>Western</u> (Krumlik and Kimmins, 1973)					
..... cont'd					
Branches, 0.6 - 2.5 cm	ODW-kg	$Y = 8.504 + 7.119 D_3 CL_m$	8	.76	15.0 - 49.0 (cm)
Branches, < 0.6 cm & foliage	ODW-kg	$Y = -2.601 + 7.480 D_3 CL_m$	8	.88	15.0 - 49.0 (cm)
Branches, < 0.6 cm	ODW-kg	$Y = 0.772 + 2.296 D_3 CL_m$	8	.79	15.0 - 49.0 (cm)
Foliage	ODW-kg	$Y = -3.340 + 5.176 D_3 CL_m$	8	.88	15.0 - 49.0 (cm)
Tree, a.s. 30 cm	ODW-kg	$\log Y = 2.304 + 0.850 \log D_7$	8	.99	15.0 - 49.0 (cm)
Stem, a.s. 30 cm, wood	ODW-kg	$\log Y = 2.112 + 0.870 \log D_7$	8	.99	15.0 - 49.0 (cm)
Stem, a.s. 30 cm, bark	ODW-kg	$\log Y = 3.060 + 1.197 \log BA$	8	.94	15.0 - 49.0 (cm)
Branches, > 2.5 cm	ODW-kg	$\log Y = 3.270 + 3.868 \log D_3$	8	.78	15.0 - 49.0 (cm)
Branches, 0.6 - 2.5 cm	ODW-kg	$\log Y = 0.936 + 1.007 \log D_3 CL_m$	8	.91	15.0 - 49.0 (cm)
Branches, < 0.6 cm & foliage	ODW-kg	$\log Y = 0.628 + 1.272 \log D_3 CL_m$	8	.96	15.0 - 49.0 (cm)
Branches, < 0.6 cm	ODW-kg	$\log Y = -0.213 + 0.697 \log D_3 (CL_m)^2$	8	.80	15.0 - 49.0 (cm)
Foliage	ODW-kg	$\log Y = -0.816 + 1.114 \log D_3 (CL_m)^2$	8	.84	15.0 - 49.0 (cm)
<u>Hemlock</u>					
<u>Western</u> (Kurucz, 1961)					
Tree	ODW-lbs	$Y = 0.071955 D_6$	89	.98	2.5 - 90 (cm)
Stem - wood	ODW-lbs	$Y = 0.056537 D_6$	89	.99	2.5 - 90 (cm)
Stem - bark	ODW-lbs	$Y = 0.008406 D_6$	89	.93	2.5 - 90 (cm)
Crown	ODW-lbs	$Y = 0.165095 DC_1 + 0.000255 DC_2$	89	.81	2.5 - 90 (cm)
Branches > 3/4 in	ODW-lbs	$Y = -0.052963 DC_1 + 0.000203 DC_2$	89	.82	2.5 - 90 (cm)
Branches 1/4 - 3/4 in	ODW-lbs	$Y = 0.039065 DC_1 + 0.000015 DC_2$	89	.80	2.5 - 90 (cm)
Branches < 1/4 in	ODW-lbs	$Y = 0.047008 DC_1 + 0.000019 DC_2$	89	.79	2.5 - 90 (cm)
Foliage	ODW-lbs	$Y = 0.131984 DC_1 + 0.000018 DC_2$	89	.75	2.5 - 90 (cm)
<u>Hobblebush</u>					
<u>(Teifer, 1969)</u>					
Tree	ODW-g	$\ln Y = -4.079 + 3.243 \ln D_4$	19	.95	2.8 - 15.9 (cm)
Foliage	ODW-g	$\ln Y = -4.347 + 2.679 \ln D_4$	19	.93	2.8 - 15.9 (cm)

Appendix

SPECIES (Reference) Equation Title	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Honeysuckle</u> American Fly (Telfer, 1969)					
Tree	ODW-g	$\ln Y = -2.427 + 2.770 \ln D_4$	20	.96	1.2 — 6.9 (mm)
Foliage	ODW-g	$\ln Y = -2.776 + 2.398 \ln D_4$	20	.86	1.2 — 6.9 (mm)
<u>Holly</u> Canada (Telfer, 1969)					
Tree	ODW-g	$\ln Y = -3.711 + 3.340 \ln D_4$	20	.94	1.9 — 9.0 (mm)
Foliage	ODW-g	$\ln Y = -4.190 + 2.851 \ln D_4$	20	.93	1.9 — 9.0 (mm)
<u>Holly</u> False (Telfer, 1969)					
Tree	ODW-g	$\ln Y = -3.040 + 2.819 \ln D_4$	20	.97	1.6 — 28.5 (mm)
Foliage	ODW-g	$\ln Y = -3.691 + 2.231 \ln D_4$	20	.97	1.6 — 28.5 (mm)
<u>Huckleberry</u> Black (Telfer, 1969)					
Tree	ODW-g	$\ln Y = -1.159 + 2.156 \ln D_4$	22	.86	4.6 — 25.7 (mm)
Foliage	ODW-g	$\ln Y = -0.823 + 1.502 \ln D_4$	22	.65	4.6 — 25.7 (mm)
<u>Huckleberry</u> Black (Whittaker and Woodwell, 1968)					
Stem	ODW-g	$\log Y = 1.3681 + 3.1008 \ln D_5$	15	.96 (r)	No data available
Stem wood	ODW-g	$\log Y = 1.2449 + 3.1767 \ln D_5$	15	.95 (r)	No data available
Stem bark	ODW-g	$\log Y = 0.6536 + 2.6971 \ln D_5$	15	.89 (r)	No data available
Branches	ODW-g	$\log Y = 1.1167 + 3.0648 \ln D_5$	15	.90 (r)	No data available
Tree	ODW-g	$\log Y = 2.4200 + 1.8627 \ln D_5$	15	.96 (r)	No data available
<u>Inkberry</u> (Telfer, 1969)					
Tree	ODW-g	$\ln Y = -2.924 + 3.011 \ln D_4$	20	.91	2.3 — 18.7 (mm)
Foliage	ODW-g	$\ln Y = -3.326 + 2.722 \ln D_4$	20	.81	2.3 — 18.7 (mm)

Appendix

SPECIES (Reference) Equation Title	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Labrador Tea</u> (Telfer, 1969)					
Tree	ODW-g	$\ln Y = -2.894 + 2.832 \ln D_4$	22	.96	2.3 - 8.3 (mm)
Foliage	ODW-g	$\ln Y = -3.044 + 2.413 \ln D_4$	22	.90	2.3 - 8.3 (mm)
<u>Laurel</u> Sheep (Telfer, 1969)					
Tree	ODW-g	$\ln Y = -2.205 + 2.384 \ln D_4$	20	.95	0.9 - 8.7 (cm)
Foliage	ODW-g	$\ln Y = -2.675 + 2.091 \ln D_4$	20	.90	0.9 - 8.7 (cm)
<u>Leatherleaf</u> (Telfer, 1969)					
Tree	ODW-g	$\ln Y = -2.325 + 2.626 \ln D_4$	20	.94	1.4 - 8.2 (mm)
Foliage	ODW-g	$\ln Y = -2.755 + 2.220 \ln D_4$	20	.89	1.4 - 8.2 (mm)
<u>Maple</u> Red (Ribe, 1973)					
Foliage	FW-g	$\log Y = 2.4872 + 1.9012 \ln D_2$	30	.79	2.5 - 15.0 (cm)
Branches	FW-g	$\log Y = 2.5803 + 1.8741 \ln D_2$	30	.91	2.5 - 15.0 (cm)
Stem	FW-g	$\log Y = 3.1020 + 2.5855 \ln D_2$	30	.99	2.5 - 15.0 (cm)
Foliage	ODW-g	$\log Y = 2.1237 + 1.8015 \log D_2$	30	.85	2.5 - 15.0 (cm)
Branches	ODW-g	$\log Y = 2.3088 + 1.9148 \log D_2$	30	.90	2.5 - 15.0 (cm)
Stem	ODW-g	$\log Y = 2.8479 + 2.6522 \log D_2$	30	.99	2.5 - 15.0 (cm)
<u>Maple</u> Red (Telfer, 1969)					
Tree	ODW-g	$\ln Y = -4.194 + 2.094 \ln D_4$	20	.98	3.8 - 40.1 (mm)
Foliage	ODW-g	$\ln Y = -2.880 + 2.079 \ln D_4$	20	.95	3.8 - 40.1 (mm)
<u>Maple</u> Red (Young et al., 1964)					
Tree tot.	FW-lbs	$\ln Y = .6654 + 2.2234 \ln D_1 + .3390 \ln H_1$	20	.97	15.3 - 29.0 (cm)
Tree tot.	ODW-lbs	46% of FW from above equation	20	.97	15.3 - 29.0 (cm)

Appendix

SPECIES (Reference) Equation Title	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Maple</u>					
<u>Red</u>					
(Young et al., 1964)					
... cont'd					
Tree tot. - Roots < 4"	FW-lbs	$\ln Y = .6020 + 1.9845 \ln D_1 + .4597 \ln H_1$	20	.97	15.3 - 29.0 (cm)
Tree tot. - Roots < 4"	ODW-lbs	48% of FW from above equation	20	.97	15.3 - 29.0 (cm)
Tree a.s. 6"	FW-lbs	$\ln Y = .1651 + 2.1679 \ln D_1 + .4292 \ln H_1$	20	.96	15.3 - 29.0 (cm)
Tree a.s. 6"	ODW-lbs	47% of FW from above equation	20	.96	15.3 - 29.0 (cm)
Stem a.s. 6"	FW-lbs	$\ln Y = -1.4801 + 1.8801 \ln D_1 + .9332 \ln H_1$	20	.79	15.3 - 29.0 (cm)
Stem a.s. 6"	ODW-lbs	53% of FW from above equation	20	.79	15.3 - 29.0 (cm)
Bole a.s. 6", t. 4"	FW-lbs	$\ln Y = -2.0214 + 2.0215 \ln D_1 + .9670 \ln H_1$	20	.98	15.3 - 29.0 (cm)
Bole a.s. 6", t. 4"	ODW-lbs	53% of FW from above equation	20	.98	15.3 - 29.0 (cm)
All branches	FW-lbs	$\ln Y = 1.5120 + 1.7383 \ln D_1 - .0957 \ln H_1$	20	.57	15.3 - 29.0 (cm)
All branches	ODW-lbs	29% of FW from above equation	20	.57	15.3 - 29.0 (cm)
Branches, < 1 in	FW-lbs	$\ln Y = 1.5749 + .8616 \ln D_1 + .2486 \ln H_1$	20	.29	15.3 - 29.0 (cm)
Branches, < 1 in	ODW-lbs	16% of FW from above equation	20	.29	15.3 - 29.0 (cm)
Stump 6" and roots > 1 in	FW-lbs	$\ln Y = .0730 + 2.4302 \ln D_1 - .0465 \ln H_1$	20	.84	15.3 - 29.0 (cm)
Stump 6" and roots > 1 in	ODW-lbs	49% of FW from above equation	20	.84	15.3 - 29.0 (cm)
Roots > 1 in	FW-lbs	$\ln Y = .4463 + 2.2539 \ln D_1 - .1634 \ln H_1$	20	.70	15.3 - 29.0 (cm)
Roots > 1 in	ODW-lbs	47% of FW from above equation	20	.70	15.3 - 29.0 (cm)
Roots < 1 in	FW-lbs	$\ln Y = -1.4358 + 1.6306 \ln D_1 + .3564 \ln H_1$	20	.70	15.3 - 29.0 (cm)
Roots < 1 in	ODW-lbs	27% of FW from above equation	20	.70	15.3 - 29.0 (cm)
<u>Maple</u>					
<u>Red</u>					
(Young and Carpenter, 1967)					
Foliage	FW-g	$\ln Y = 0.770 + 1.917 \ln H_1$	40	.90	2.5 - 10.0 (cm)
Branches	FW-g	$\ln Y = -1.007 + 2.594 \ln H_1$	40	.88	2.5 - 10.0 (cm)
Stem	FW-g	$\ln Y = -0.221 + 2.841 \ln H_1$	40	.97	2.5 - 10.0 (cm)
Roots	FW-g	$\ln Y = 0.196 + 2.473 \ln H_1$	40	.96	2.5 - 10.0 (cm)
Tree tot.	FW-g	$\ln Y = 1.505 + 2.479 \ln H_1$	40	.95	2.5 - 10.0 (cm)

Appendix

SPECIES (Reference) Equation Title	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Maple</u> Red (Young and Carpenter, 1967) cont'd					
Foliage	ODW-g	$\ln Y = -0.217 + 1.923 \ln H_1$	40	.87	2.5 — 10.0 (cm)
Branches	ODW-g	$\ln Y = -1.583 + 2.577 \ln H_1$	40	.87	2.5 — 10.0 (cm)
Stem	ODW-g	$\ln Y = -0.789 + 2.842 \ln H_1$	40	.96	2.5 — 10.0 (cm)
Roots	ODW-g	$\ln Y = -0.538 + 2.497 \ln H_1$	40	.94	2.5 — 10.0 (cm)
Tree tot.	ODW-g	$\ln Y = +0.728 + 2.532 \ln H_1$	40	.95	2.5 — 10.0 (cm)
<u>Maple</u> Striped (Telfer, 1969)					
Tree	ODW-g	$\ln Y = -3.518 + 2.878 \ln D_4$	20	.99	9.2 — 43.1 (mm)
Foliage	ODW-g	$\ln Y = -3.334 + 2.220 \ln D_4$	20	.98	9.2 — 43.1 (mm)
<u>Maple</u> Sugar (Ribe, 1973)					
Foliage	FW-g	$\log Y = 2.4060 + 1.6400 \log D_2$	30	.83	2.5 — 15.0 (cm)
Branches	FW-g	$\log Y = 2.6141 + 1.5525 \log D_2$	30	.85	2.5 — 15.0 (cm)
Stem	FW-g	$\log Y = 3.2384 + 2.4834 \log D_2$	30	.99	2.5 — 15.0 (cm)
Foliage	ODW-g	$\log Y = 2.0383 + 1.6701 \log D_2$	30	.83	2.5 — 15.0 (cm)
Branches	ODW-g	$\log Y = 2.4004 + 1.5571 \log D_2$	30	.85	2.5 — 15.0 (cm)
Stem	ODW-g	$\log Y = 3.0609 + 2.4927 \log D_2$	30	.99	2.5 — 15.0 (cm)
<u>Oak</u> Black (King and Schnell, 1972)					
Tree tot., wood	FW-lbs	$\log Y = 1.12282 + 2.12073 \log D_1$	26	.98	28.8 — 87.0 (cm)
Tree tot., bark	FW-lbs	$\log Y = 0.53961 + 2.01311 \log D_1$	26	.95	28.8 — 87.0 (cm)
Tree tot.	FW-lbs	$\log Y = 1.22080 + 2.10416 \log D_1$	26	.98	28.8 — 87.0 (cm)
Tree tot., wood	ODW-lbs	$\log Y = 0.90951 + 2.11178 \log D_1$	26	.97	28.8 — 87.0 (cm)
Tree tot., bark	ODW-lbs	$\log Y = 0.27002 + 2.08383 \log D_1$	26	.96	28.8 — 87.0 (cm)

Appendix

SPECIES (Reference)	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST-HEIGHT IN CM
<u>Oak</u>					
Black					
(King and Schnell, 1972)					
.... cont'd					
Tree tot.	ODW-lbs	$\log Y = 1.00005 + 2.10621 \log D_1$	26	.97	28.8 - 87.0 (cm)
Branches - wood	FW-lbs	$\log Y = 0.53868 + 2.09742 \log D_1$	26	.88	28.8 - 87.0 (cm)
Branches - bark	FW-lbs	$\log Y = 0.08437 + 2.08487 \log D_1$	26	.86	28.8 - 87.0 (cm)
Branches	FW-lbs	$\log Y = 0.70890 + 2.09360 \log D_1$	26	.88	28.8 - 87.0 (cm)
Branches - wood	ODW-lbs	$\log Y = 0.42075 + 2.06105 \log D_1$	26	.87	28.8 - 87.0 (cm)
Branches - bark	ODW-lbs	$\log Y = -0.22974 + 2.19873 \log D_1$	26	.88	28.8 - 87.0 (cm)
Branches	ODW-lbs	$\log Y = 0.50580 + 2.09357 \log D_1$	26	.88	28.8 - 87.0 (cm)
Bole, s 12", t 8" i.b. - wood	FW-lbs	$\log Y = 0.79041 + 2.13126 \log D_1$	26	.96	28.8 - 87.0 (cm)
Bole, s 12", t > 8" i.b. - bark	FW-lbs	$\log Y = 0.22951 + 1.77026 \log D_1$	26	.84	28.8 - 87.0 (cm)
Bole, s 12", t > 8" i.b.	FW-lbs	$\log Y = 0.86878 + 2.10122 \log D_1$	26	.95	28.8 - 87.0 (cm)
Bole, s 12", t > 8" i.b. - wood	ODW-lbs	$\log Y = 0.55932 + 2.12847 \log D_1$	26	.95	28.8 - 87.0 (cm)
Bole, s 12", t > 8" i.b. - bark	ODW-lbs	$\log Y = -0.02290 + 1.84814 \log D_1$	26	.84	28.8 - 87.0 (cm)
Bole, s 12", t > 8" i.b.	ODW-lbs	$\log Y = 0.63832 + 2.10813 \log D_1$	26	.93	28.8 - 87.0 (cm)
Stump, s 12" and root - wood	FW-lbs	$\log Y = 0.51989 + 2.10556 \log D_1$	26	.99	28.8 - 87.0 (cm)
Stump, s 12" and root - bark	FW-lbs	$\log Y = -0.07786 + 2.10255 \log D_1$	26	.99	28.8 - 87.0 (cm)
Stump, s 12" and root	FW-lbs	$\log Y = 0.61783 + 2.10475 \log D_1$	26	.99	28.8 - 87.0 (cm)
Stump, s 12" and root - wood	ODW-lbs	$\log Y = 0.28238 + 2.12145 \log D_1$	26	.99	28.8 - 87.0 (cm)
Stump, s 12" and root - bark	ODW-lbs	$\log Y = -0.31613 + 2.11888 \log D_1$	26	.99	28.8 - 87.0 (cm)
Stump, s 12" and root	ODW-lbs	$\log Y = 0.38000 + 2.12094 \log D_1$	26	.99	28.8 - 87.0 (cm)
Bole, s 12", t > 8" i.b. and branches - wood	FW-lbs	$\log Y = 0.99826 + 2.12553 \log D_1$	26	.98	28.8 - 87.0 (cm)
Bole, s 12", t > 8" i.b. and branches - bark	FW-lbs	$\log Y = 0.42651 + 1.97309 \log D_1$	26	.94	28.8 - 87.0 (cm)
Bole, s 12", t > 8" i.b. and branches	FW-lbs	$\log Y = 1.09501 + 2.10479 \log D_1$	26	.96	28.8 - 87.0 (cm)
Bole, s 12", t > 8" i.b. and branches - wood	ODW-lbs	$\log Y = 0.79251 + 2.10880 \log D_1$	26	.97	28.8 - 87.0 (cm)

Appendix

SPECIES (Reference) Equation Title	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Oak</u>					
<u>Black</u>					
(King and Schnell, 1972)					
.... cont'd					
Bole, s 12", t > 8" i.b. and branches - bark	ODW-lbs	$\log Y = 0.13817 + 2.07142 \log D_1$	26	.95	28.8 - 87.0 (cm)
Bole, s 12", t > 8" i.b. and branches	ODW-lbs	$\log Y = 0.88191 + 2.10156 \log D_1$	26	.97	28.8 - 87.0 (cm)
Branches, > 8" - wood	FW-lbs	$\log Y = -0.14531 + 3.24591 \log D_1$	26	.77	28.8 - 87.0 (cm)
Branches, > 8" - bark	FW-lbs	$\log Y = -0.21051 + 3.26054 \log D_1$	26	.79	28.8 - 87.0 (cm)
Branches, > 8"	FW-lbs	$\log Y = -1.136551 + 3.24902 \log D_1$	26	.78	28.8 - 87.0 (cm)
Branches, > 8" - wood	ODW-lbs	$\log Y = -1.163625 + 3.19694 \log D_1$	26	.72	28.8 - 87.0 (cm)
Branches, > 8" - bark	ODW-lbs	$\log Y = -2.37246 + 3.34949 \log D_1$	26	.80	28.8 - 87.0 (cm)
Branches, > 8"	ODW-lbs	$\log Y = -1.57207 + 3.23439 \log D_1$	26	.75	28.8 - 87.0 (cm)
Branches 4" to 8" - wood	FW-lbs	$\log Y = 0.26221 + 1.90836 \log D_1$	26	.69	28.8 - 87.0 (cm)
Branches 4" to 8" - bark	FW-lbs	$\log Y = -0.56453 + 2.13045 \log D_1$	26	.76	28.8 - 87.0 (cm)
Branches 4" to 8"	FW-lbs	$\log Y = 0.30818 + 1.95884 \log D_1$	26	.71	28.8 - 87.0 (cm)
Branches 4" to 8" - wood	ODW-lbs	$\log Y = 0.044075 + 1.90711 \log D_1$	26	.78	28.8 - 87.0 (cm)
Branches 4" to 8" - bark	ODW-lbs	$\log Y = -0.84576 + 2.22581 \log D_1$	26	.78	28.8 - 87.0 (cm)
Branches 4" to 8"	ODW-lbs	$\log Y = -0.06631 + 1.98757 \log D_1$	26	.72	28.8 - 87.0 (cm)
Branches, > 4" - wood	FW-lbs	$\log Y = -0.17751 + 2.51058 \log D_1$	26	.85	28.8 - 87.0 (cm)
Branches, > 4" - bark	FW-lbs	$\log Y = -0.86033 + 2.58256 \log D_1$	26	.87	28.8 - 87.0 (cm)
Branches, > 4"	FW-lbs	$\log Y = -0.09636 + 2.52525 \log D_1$	26	.86	28.8 - 87.0 (cm)
Branches, > 4" - wood	ODW-lbs	$\log Y = -0.37233 + 2.48400 \log D_1$	26	.84	28.8 - 87.0 (cm)
Branches, > 4" - bark	ODW-lbs	$\log Y = -0.11328 + 2.67315 \log D_1$	26	.88	28.8 - 87.0 (cm)
Branches, > 4"	ODW-lbs	$\log Y = -0.31213 + 2.52763 \log D_1$	26	.86	28.8 - 87.0 (cm)
Branches 2" to 4" - wood	FW-lbs	$\log Y = 0.85390 + 1.29387 \log D_1$	26	.51	28.8 - 87.0 (cm)
Branches 2" to 4" - bark	FW-lbs	$\log Y = 0.05307 + 1.54960 \log D_1$	26	.57	28.8 - 87.0 (cm)
Branches 2" to 4"	FW-lbs	$\log Y = 0.90694 + 1.35239 \log D_1$	26	.54	28.8 - 87.0 (cm)
Branches 2" to 4" - wood	ODW-lbs	$\log Y = 0.66439 + 1.27907 \log D_1$	26	.50	28.8 - 87.0 (cm)

Appendix

SPECIES (Reference)	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
Oak					
Black (King and Schnell, 1972)					
... cont'd					
Branches 2" to 4" - bark	ODW-lbs	$\log Y = -0.25166 + 1.65302 \log D_1$	26	.60	28.8 - 87.0 (cm)
Branches 2" to 4"	ODW-lbs	$\log Y = 0.68104 + 1.37377 \log D_1$	26	.57	28.8 - 87.0 (cm)
Branches, > 2" - wood	FW-lbs	$\log Y = 0.27107 + 2.25755 \log D_1$	26	.96	28.8 - 87.0 (cm)
Branches, > 2" - bark	FW-lbs	$\log Y = -0.36158 + 2.31293 \log D_1$	26	.96	28.8 - 87.0 (cm)
Branches, > 2"	FW-lbs	$\log Y = 0.36445 + 2.26755 \log D_1$	26	.97	28.8 - 87.0 (cm)
Branches, > 2" - wood	ODW-lbs	$\log Y = -0.09138 + 2.22281 \log D_1$	26	.96	28.8 - 87.0 (cm)
Branches, > 2" - bark	ODW-lbs	$\log Y = -0.65505 + 2.41620 \log D_1$	26	.97	28.8 - 87.0 (cm)
Branches, > 2"	ODW-lbs	$\log Y = 0.15585 + 2.26683 \log D_1$	26	.96	28.8 - 87.0 (cm)
Branches 1" to 2" - wood	FW-lbs	$\log Y = 0.41738 + 1.36260 \log D_1$	26	.58	28.8 - 87.0 (cm)
Branches 1" to 2" - bark	FW-lbs	$\log Y = -0.21154 + 1.48854 \log D_1$	26	.56	28.8 - 87.0 (cm)
Branches 1" to 2"	FW-lbs	$\log Y = 0.50234 + 1.39658 \log D_1$	26	.59	28.8 - 87.0 (cm)
Branches 1" to 2" - wood	ODW-lbs	$\log Y = 0.24444 + 1.34217 \log D_1$	26	.58	28.8 - 87.0 (cm)
Branches 1" to 2" - bark	ODW-lbs	$\log Y = -0.50480 + 1.56628 \log D_1$	26	.60	28.8 - 87.0 (cm)
Branches 1" to 2"	ODW-lbs	$\log Y = 0.30079 + 1.39979 \log D_1$	26	.59	28.8 - 87.0 (cm)
Branches 1/2" to 1" - wood	FW-lbs	$\log Y = 0.11432 + 1.45778 \log D_1$	26	.49	28.8 - 87.0 (cm)
Branches 1/2" to 1" - bark	FW-lbs	$\log Y = -0.16487 + 1.38758 \log D_1$	26	.51	28.8 - 87.0 (cm)
Branches 1/2" to 1"	FW-lbs	$\log Y = 0.30002 + 1.43540 \log D_1$	26	.51	28.8 - 87.0 (cm)
Branches 1/2" to 1" - wood	ODW-lbs	$\log Y = -0.06862 + 1.44493 \log D_1$	26	.49	28.8 - 87.0 (cm)
Branches 1/2" to 1" - bark	ODW-lbs	$\log Y = -0.47010 + 1.46578 \log D_1$	26	.55	28.8 - 87.0 (cm)
Branches 1/2" to 1"	ODW-lbs	$\log Y = 0.07851 + 1.45131 \log D_1$	26	.52	28.8 - 87.0 (cm)
Branches < 1/2" - wood	FW-lbs	$\log Y = 0.06130 + 1.34446 \log D_1$	26	.53	28.8 - 87.0 (cm)
Branches, < 1/2" - bark	FW-lbs	$\log Y = -0.25690 + 1.41209 \log D_1$	26	.50	28.8 - 87.0 (cm)
Branches, < 1/2"	FW-lbs	$\log Y = 0.23577 + 1.36564 \log D_1$	26	.53	28.8 - 87.0 (cm)
Branches, < 1/2" - wood	ODW-lbs	$\log Y = -0.10678 + 1.31634 \log D_1$	26	.53	28.8 - 87.0 (cm)
Branches, < 1/2" - bark	ODW-lbs	$\log Y = -0.59374 + 1.51130 \log D_1$	26	.56	28.8 - 87.0 (cm)

Appendix

SPECIES (Reference)	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM/OR DIAMETER AT BREAST HEIGHT IN CM
<u>Oak</u>					
<u>Black</u>					
(King and Schnell, 1972)					
..... cont'd					
Branches, < 1/2"	ODW-lbs	$\log Y = 0.00817 + 1.38250 \log D_1$	26	.56	28.8 - 87.0 (cm)
Branches, < 2" - wood	FW-lbs	$\log Y = 0.68607 + 1.40754 \log D_1$	26	.62	28.8 - 87.0 (cm)
Branches, < 2" - bark	FW-lbs	$\log Y = 0.25774 + 1.44729 \log D_1$	26	.63	28.8 - 87.0 (cm)
Branches, < 2"	FW-lbs	$\log Y = 0.82284 + 1.42063 \log D_1$	26	.63	28.8 - 87.0 (cm)
Branches, < 2" - wood	ODW-lbs	$\log Y = 0.51562 + 1.38441 \log D_1$	26	.62	28.8 - 87.0 (cm)
Branches, < 2" - bark	ODW-lbs	$\log Y = -0.04710 + 1.52719 \log D_1$	26	.66	28.8 - 87.0 (cm)
Branches, < 2"	ODW-lbs	$\log Y = 0.61270 + 1.42803 \log D_1$	26	.64	28.8 - 87.0 (cm)
Branches, < 4" - wood	FW-lbs	$\log Y = 1.10606 + 1.32800 \log D_1$	26	.61	28.8 - 87.0 (cm)
Branches, < 4" - bark	FW-lbs	$\log Y = 0.48070 + 1.48380 \log D_1$	26	.63	28.8 - 87.0 (cm)
Branches, < 4"	FW-lbs	$\log Y = 1.19125 + 1.37055 \log D_1$	26	.63	28.8 - 87.0 (cm)
Branches, < 4" - wood	ODW-lbs	$\log Y = 0.92513 + 1.31021 \log D_1$	26	.61	28.8 - 87.0 (cm)
Branches, < 4" - bark	ODW-lbs	$\log Y = 0.16728 + 1.58130 \log D_1$	26	.66	28.8 - 87.0 (cm)
Branches, < 4"	ODW-lbs	$\log Y = .97158 + 1.38644 \log D_1$	26	.64	28.8 - 87.0 (cm)
Dead branches - wood	FW-lbs	$\log Y = -0.45497 + 1.79386 \log D_1$	26	.38	28.8 - 87.0 (cm)
Dead branches - bark	FW-lbs	$\log Y = -1.05839 + 1.91374 \log D_1$	26	.45	28.8 - 87.0 (cm)
Dead branches	FW-lbs	$\log Y = -0.36597 + 1.82896 \log D_1$	26	.40	28.8 - 87.0 (cm)
Dead branches - wood	ODW-lbs	$\log Y = -0.68961 + 1.87904 \log D_1$	26	.41	28.8 - 87.0 (cm)
Dead branches - bark	ODW-lbs	$\log Y = -1.26380 + 1.98905 \log D_1$	26	.50	28.8 - 87.0 (cm)
Dead branches	ODW-lbs	$\log Y = -0.58932 + 1.90855 \log D_1$	26	.43	28.8 - 87.0 (cm)
<u>Red</u>					
(Telfer, 1969)					
Tree	ODW-g	$\ln Y = -2.299 + 2.649 \ln D_4$	20	.99	2.2 - 39.9 (cm)
Foliage	ODW-g	$\ln Y = -2.602 + 2.198 \ln D_4$	20	.97	2.2 - 39.9 (cm)

Appendix

SPECIES (Reference) Equation Title	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Oak</u>					
<u>Scarlet</u> (Whittaker and Woodwell, 1968)					
Branches	ODW-g	$\log Y = 1.0745 + 2.889 \log D_2$	15	.97	No data available
Tree	ODW-g	$\log Y = 2.3948 + 2.1900 \log D_2$	15	.99	No data available
Stem	ODW-g	$\log Y = 2.1564 + 2.2391 \log D_2$	15	.99	No data available
Stem wood	ODW-g	$\log Y = 2.0001 + 2.3025 \log D_2$	15	.99	No data available
Stem bark	ODW-g	$\log Y = 1.6767 + 1.9909 \log D_2$	15	.99	No data available
Stem heart wood	ODW-g	$\log Y = 1.8955 + 2.0316 \log D_2$	15	.99	No data available
<u>Oak</u>					
<u>Scrub</u> (Whittaker and Woodwell, 1968)					
Stem	ODW-g	$\log Y = 1.3561 + 2.7146 \log D_5$	15	.97	0 - 3 (cm)
Stem wood	ODW-g	$\log Y = 1.2442 + 2.7663 \log D_5$	15	.97	0 - 3 (cm)
Stem bark	ODW-g	$\log Y = 0.6958 + 2.5831 \log D_5$	15	.97	0 - 3 (cm)
Branches	ODW-g	$\log Y = 0.8596 + 3.2040 \log D_5$	15	.92	0 - 3 (cm)
Tree	ODW-g	$\log Y = 1.7076 + 2.3551 \log D_5$	15	.95	0 - 3 (cm)
<u>Oak</u>					
<u>White</u> (Whittaker and Woodwell, 1968)					
Stem	ODW-g	$\log Y = 2.1436 + 2.1844 \log D_2$	No Data	.99	No data available
Stem wood	ODW-g	$\log Y = 1.9610 + 2.2537 \log D_2$	No Data	.99	No data available
Stem bark	ODW-g	$\log Y = 1.7075 + 1.9747 \log D_2$	No Data	.99	No data available
Stem heart wood	ODW-g	$\log Y = 1.3220 + 2.4628 \log D_2$	No Data	.96	No data available
Branches	ODW-g	$\log Y = 1.4383 + 2.3371 \log D_2$	No Data	.96	No data available
Tree	ODW-g	$\log Y = 2.3058 + 2.1666 \log D_2$	No Data	.99	No data available
<u>Pine</u>					
<u>Jack</u> (Brown, 1965) Live & dead slash, t. < 1.5"	ODW-lbs	$\log Y = 0.8234 + 0.133 \log D_1 + 0.003393 \log S1$ $- 0.0001127 \log TA$	95	.90	5.0 - 25.0 (cm)

Appendix

SPECIES (Reference)	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Pine</u> Jack (Brown, 1965) cont'd					
Live slash, t. < 1.5"	ODW-lbs	$\log Y = 0.9733 + 0.1148 \log D_1 - 0.0001556 \log TA$	95	.82	5.0 — 25.0 (cm)
Dead branches	ODW-lbs	$\log Y = 0.2087 \log D_1 - 0.0978$	95	.78	5.0 — 25.0 (cm)
<u>Pine</u> Jack (good site) (Doucet, et al., 1976)					
Stem wood	ODW-g	$\log Y = 0.847345 + 1.430786 \log D_2 + 0.827688 \log H_3$	18	.99	5.3 — 21.4 (cm)
Stem bark	ODW-g	$\log Y = 0.708473 + 1.845331 \log D_2 + 0.707754 \log H_3$	18	.94	5.3 — 21.4 (cm)
Live branches	ODW-g	$\log Y = 0.525747 + 2.574138 \log D_2 + 0.210360 \log H_3$	18	.91	5.3 — 21.4 (cm)
Dead branches	ODW-g	$\log Y = -2.703526 + 0.187033 \log D_2 + 5.083131 \log H_3$	17	.90	5.3 — 21.4 (cm)
Foliage	ODW-g	$\log Y = -0.043127 + 2.106884 \log D_2 + 0.897754 \log H_3$	18	.96	5.3 — 21.4 (cm)
Cones	ODW-g	$\log Y = -1.304145 + 2.354415 \log D_2 + 0.609672 \log H_3$	18	.43	5.3 — 21.4 (cm)
Tree	ODW-g	$\log Y = 1.053679 + 1.636275 \log D_2 + 1.568117 \log H_3$	17	.99	5.3 — 21.4 (cm)
<u>Pine</u> Jack (medium site) (Doucet, et al., 1976)					
Stem wood	ODW-g	$\log Y = 1.348119 + 2.052095 \log D_2 + 0.793679 \log H_3$	18	.99	1.8 — 19.3 (cm)
Stem bark	ODW-g	$\log Y = 1.168162 + 1.852994 \log D_2 + 0.306818 \log H_3$	18	.97	1.8 — 19.3 (cm)
Live branches	ODW-g	$\log Y = 1.237128 + 4.539180 \log D_2 - 2.280269 \log H_3$	18	.99	1.8 — 19.3 (cm)
Dead branches	ODW-g	$\log Y = 1.294463 + 2.826062 \log D_2 - 0.943957 \log H_3$	18	.88	1.8 — 19.3 (cm)
Foliage	ODW-g	$\log Y = 0.077329 + 4.008232 \log D_2 - 0.914902 \log H_3$	18	.96	1.8 — 19.3 (cm)
Cones	ODW-g	$\log Y = -4.158016 + 3.553876 \log D_2 + 2.396761 \log H_3$	16	.45	1.8 — 19.3 (cm)
Tree	ODW-g	$\log Y = 1.752336 + 2.447128 \log D_2 + 0.196253 \log H_3$	18	.45	1.8 — 19.3 (cm)
<u>Pine</u> Jack (Hegyí, 1972)					
Tree	ODW-kg	$\log Y = -1.0368 + 2.4206 \log D_2$	77	.99	2.9 — 31.7 (cm)
Stem	ODW-kg	$\log Y = -1.5481 + 2.7144 \log D_2$	77	.95	2.9 — 31.7 (cm)
Crown	ODW-kg	$\log Y = -1.1810 + 1.9222 \log D_2$	77	.77	2.9 — 31.7 (cm)

Appendix

SPECIES (Reference). Equation Title	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Pine</u>					
<u>Jack</u>					
(Hegvi, 1972)					
... cont'd					
Lower 1/3 of crown	ODW-kg	$\log Y = -1.4178 + 1.7458 \log D_2$	77	.65	2.9 - 31.7 (cm)
Middle 1/3 of crown	ODW-kg	$\log Y = -1.6155 + 1.9968 \log D_2$	77	.75	2.9 - 31.7 (cm)
Upper 1/3 of crown	ODW-kg	$\log Y = -2.2699 + 2.1090 \log D_2$	77	.83	2.9 - 31.7 (cm)
Foliage	ODW-kg	$\log Y = -1.4289 + 1.7188 \log D_2$	77	.70	2.9 - 31.7 (cm)
Cones	ODW-kg	$\log Y = -4.9319 + 3.8667 \log D_2$	77	.72	2.9 - 31.7 (cm)
<u>Pine</u>					
<u>Jack</u>					
(Keen, 1963)					
Bole, s*, t 3"	FW-lbs	$\log Y = 0.611 + 2.301 \log D_1$	89	.98	10.0 - 40.0 (cm)
Tree, s*	FW-lbs	$\log Y = 0.777 + 2.214 \log D_1$	89	.98	10.0 - 40.0 (cm)
Bole, s*, t 3"	FW-lbs	$\log Y = -0.9458 + 2.060 \log D_1 + 0.9831 \log H_1$	78	.99	10.0 - 40.0 (cm)
Slash	FW-lbs	$\log Y = 0.180 + 2.039 \log D_1$	90	.70	10.0 - 40.0 (cm)
* = D ₁					
<u>Pine</u>					
<u>Lodgepole</u>					
(Gary, 1976)					
Needles & branchwood	ODW-kg	$Y = -10.594 + 1.528 D_2$	117	.78	6.6 - 19.8 (cm)
Needles & branchwood	ODW-kg	$Y = -8.727 + 1.656 D$	117	.78	6.6 - 19.8 (cm)
Needles & branchwood	ODW-kg	$Y = -3.809 + 2.070 CL$	117	.33	6.6 - 19.8 (cm)
Needles & branchwood	ODW-kg	$Y = -17.672 + 2.538 H_3$	117	.44	6.6 - 19.8 (cm)
Needles & branchwood	ODW-kg	$Y = -2.442 + 0.145 D_2 CL$	117	.68	6.6 - 19.8 (cm)
Needles & branchwood	ODW-kg	$Y = -5.204 + 0.105 D_2 H_3$	117	.74	6.6 - 19.8 (cm)
Needles & branchwood	ODW-kg	$Y = -0.833 + 0.005 D_7 H_3$	117	.80	6.6 - 19.8 (cm)
Needles & branchwood	ODW-kg	$Y = 0.605 + 0.004 (D_2)^3$	117	.83	6.6 - 19.8 (cm)
Needles & branchwood	ODW-kg	$Y = -3.300 + 0.154 D_2 + 0.061 (D_2)^2$	117	.82	6.6 - 19.8 (cm)
Needles & branchwood	ODW-kg	$Y = 0.0047 D_2^{2.916}$	117	.91	6.6 - 19.8 (cm)

Appendix

SPECIES (Reference) Equation Title	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Pine</u> Lodgepole (Johnstone, 1970)					
Tree tot.	ODW-lbs	$\log Y = -0.9664 + 0.9971 \log D_6$	85	.99	10.0 - 33.5 (cm)
Tree, s 12"	ODW-lbs	$\log Y = -0.9878 + 0.9853 \log D_6$	85	.98	10.0 - 33.5 (cm)
Stem, s 12"	ODW-lbs	$\log Y = -0.8887 + 0.9381 \log D_6$	85	.98	10.0 - 33.5 (cm)
Stem, s 12", bark	ODW-lbs	$\log Y = -2.5841 + 1.1395 \log D_6$	85	.86	10.0 - 33.5 (cm)
Needles	ODW-lbs	$\log Y = -2.9508 + 1.1480 \log D_6$	85	.85	10.0 - 33.5 (cm)
Branches	ODW-lbs	$\log Y = -4.1255 + 1.5087 \log D_6$	85	.89	10.0 - 33.5 (cm)
Stump & root	ODW-lbs	$\log Y = -1.880 + 1.022 \log D_6$	85	.95	10.0 - 33.5 (cm)
Roots	ODW-lbs	$\log Y = -2.1045 + 1.0704 \log D_6$	85	.94	10.0 - 33.5 (cm)
<u>Pine</u> Lodgepole (Kil, 1967)					
Branch - live	ODW-lbs	$\log Y = -0.67346 + 1.26470 \log D_1 + 1.36575 \log CW$	101	.95	2.3 - 52.8 (cm)
Crown	ODW-lbs	$\log Y = -0.642994 + 1.30603 \log D_1 + 1.37732 \log CW$	101	.96	2.3 - 52.8 (cm)
<u>Pine</u> Lodgepole (Kimmins, 1974)					
Bole, a.s. 30.5 cm, t. 2.5 cm, wood	ODW-kg	$\log Y = 2.405 + 1.222 \log D_7 - 0.121 \log D_1$	13	.99	10.0 - 32.0 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, bark	ODW-kg	$\log Y = 3.025 + 2.207 \log D_3$	13	.95	10.0 - 32.0 (cm)
Branches, .64 - 2.5 cm	ODW-kg	$\log Y = 0.530 + 1.101 \log D_7 + 0.399 \log D_7$	13	.97	10.0 - 32.0 (cm)
Branches, < .64 cm	ODW-kg	$\log Y = 0.271 + 1.189 \log D_3 C_{Lm}$	13	.88	10.0 - 32.0 (cm)
Foliage	ODW-kg	$\log Y = 0.562 + 1.322 \log D_7$	13	.95	10.0 - 32.0 (cm)
<u>Pine</u> Pitch (Whittaker and Woodwell, 1968)					
Stem	ODW-g	$\log Y = 1.8758 + 2.3261 \log D_2$	15	.99	No data available
Stem wood	ODW-g	$\log Y = 1.7226 + 2.3779 \log D_2$	15	.99	No data available
Stem bark	ODW-g	$\log Y = 1.3832 + 2.1249 \log D_2$	15	.99	No data available

Appendix

SPECIES (Reference)	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Pine</u> Pitch (Whittaker and Woodwell, 1968) cont'd					
Stem heart wood	ODW-g	$\log Y = 1.7553 + 2.3291 \log D_2$	15	.99	No data available
Branches	ODW-g	$\log Y = 1.1100 + 2.5516 \log D_2$	15	.98	No data available
Tree	ODW-g	$\log Y = 2.0171 + 2.3373 \log D_2$	15	.99	No data available
<u>Pine</u> Red (Brown, 1965) Live slash t. < 1.5"	ODW-lbs	$\log Y = 0.9990 + 0.0946 \log D_1 + 0.003614 \log SI - 0.0002722 \log TA$	103	.86	5.0 - 25.0 (cm)
<u>Pine</u> Scots (Hakkila, 1972) Stump* and Roots > 5 cm * s height = 4.7 + .19 diameter of stump top	ODW-kg	$Y = -4.9 + 0.044 S_1^2$	213	.92 (r)	15 - 45 (cm)
<u>Pine</u> Scots (Hakkila, 1971) Branches, wood	ODW-kg	$Y = 7.2 + 0.0505 D_2^2 - 1.39 H_3 + 0.14 D_2 BR$	240	.93 (r)	10.0 - 30.0 (cm)
Slash, t < 6 cm, wood	ODW-kg	$Y = 8.7 + 0.507 D_2^2 - 1.39 H_3 + 0.14 D_2 BR$	240	.93 (r)	10.0 - 30.0 (cm)
Foliage	ODW-kg	$Y = -9.5 + 0.54 D_2 + 0.117 CR$	240	.89 (r)	7 - 30 (cm)
Slash	ODW-kg	$Y = -0.1 + 0.155 D_2^2$	240	.90 (r)	5.0 - 25.0 (cm)
Slash	ODW-kg	$Y = -0.1 + 0.00141 D_2^2 CR$	240	.94 (r)	10 - 30 (cm)
<u>Pine</u> Slash (Curtis, 1965) Bole, s 7, t 2"	FW-lbs	$Y = -60.3 + 0.145 D_6 + 2.501 A_1$	900	.98	No data available
Bole, s 7, t 2", bark	FW-lbs	$Y = 15.7 + 0.012 D_6 + 0.619 A_1$	900	.91	No data available
Bole, s 7, t 2", wood	ODW-% of FW	$Y = -1.788 A_2 - .099 A_3 + .777$	900	.46	No data available

Appendix

SPECIES (Reference) Equation Title	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Pine</u>					
<u>White</u>					
(Young, et al., 1964)					
Tree tot.	FW-lbs	$\ln Y = .6592 + 2.2234 \ln D_1 + .3390 \ln H_1$	27	.98	14.5 — 24.5 (cm)
Tree tot.	ODW-lbs	36% of FW from above equation	27	.98	14.5 — 24.5 (cm)
Tree tot. — Roots < 4"	FW-lbs	$\ln Y = .2050 + 2.1883 \ln D_1 + .4514 \ln H_1$	27	.98	14.5 — 24.5 (cm)
Tree tot. — Roots < 4"	ODW-lbs	37% of FW from above equation	27	.98	14.5 — 24.5 (cm)
Tree a.s. 6"	FW-lbs	$\ln Y = .2150 + 2.1679 \ln D_1 + .4292 \ln H_1$	27	.98	14.5 — 24.5 (cm)
Tree a.s. 6"	ODW-lbs	36% of FW from above equation	27	.98	14.5 — 24.5 (cm)
Stem a.s. 6"	FW-lbs	$\ln Y = -1.4111 + 1.8801 \ln D_1 + .9332 \ln H_1$	27	.83	14.5 — 24.5 (cm)
Stem a.s. 6"	ODW-lbs	40% of FW from above equation	27	.83	14.5 — 24.5 (cm)
Bole a.s. 6", t. 4"	FW-lbs	$\ln Y = -1.9231 + 2.0215 \ln D_1 + .9670 \ln H_1$	27	.98	14.5 — 24.5 (cm)
Bole a.s. 6", t. 4"	ODW-lbs	40% of FW from above equation	27	.98	14.5 — 24.5 (cm)
All branches	FW-lbs	$\ln Y = 3.6340 + 3.6004 \ln D_1 - 1.6346 \ln H_1$	27	.91	14.5 — 24.5 (cm)
All branches	ODW-lbs	16% of FW from above equation	27	.91	14.5 — 24.5 (cm)
Branches < 1 in	FW-lbs	$\ln Y = 2.7899 + 2.7716 \ln D_1 - 1.0629 \ln H_1$	27	.87	14.5 — 24.5 (cm)
Branches < 1 in	ODW-lbs	13% of FW from above equation	27	.87	14.5 — 24.5 (cm)
Stump 6" and roots > 1 in	FW-lbs	$\ln Y = 2.0191 + 3.2156 \ln D_1 - .9513 \ln H_1$	27	.93	14.5 — 24.5 (cm)
Stump 6" and roots > 1 in	ODW-lbs	38% of FW from above equation	27	.93	14.5 — 24.5 (cm)
Roots > 1 in	FW-lbs	$\ln Y = 1.7421 + 3.2955 \ln D_1 - 1.0554 \ln H_1$	27	.91	14.5 — 24.5 (cm)
Roots > 1 in	ODW-lbs	38% of FW from above equation	27	.91	14.5 — 24.5 (cm)
Roots > 1 in	FW-lbs	$\ln Y = -1.8053 + 1.6306 \ln D_1 + .3564 \ln H_1$	27	.85	14.5 — 24.5 (cm)
Roots < 1 in	ODW-lbs	21% of FW from above equation	27	.85	14.5 — 24.5 (cm)
<u>Pine</u>					
<u>White sp.</u>					
(Young and Carpenter, 1967)					
Foliage	FW-g	$\ln Y = 1.663 + 1.752 \ln H_1$	10	.89	0.0 — 75 (mm)
Branches	FW-g	$\ln Y = 1.253 + 2.007 \ln H_1$	10	.94	0.0 — 75 (mm)
Stem	FW-g	$\ln Y = 0.346 + 2.913 \ln H_1$	10	.99	0.0 — 75 (mm)
Roots	FW-g	$\ln Y = 0.831 + 2.379 \ln H_1$	10	.94	0.0 — 75 (mm)

Appendix

SPECIES (Reference) Equation Title	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Pine</u>					
<u>White sp.</u>					
(Young and Carpenter, 1967)					
.... cont'd					
Tree tot.	FW-g	$\ln Y = 2.492 + 2.346 \ln H_1$	10	.95	0.0 - 75 (mm)
Foliage	ODW-g	$\ln Y = 0.808 + 1.750 \ln H_1$	10	.89	0.0 - 75 (mm)
Branches	ODW-g	$\ln Y = 0.413 + 2.064 \ln H_1$	10	.95	0.0 - 75 (mm)
Stem	ODW-g	$\ln Y = -0.177 + 2.814 \ln H_1$	10	.98	0.0 - 75 (mm)
Roots	ODW-g	$\ln Y = -0.081 + 2.401 \ln H_1$	10	.94	0.0 - 75 (mm)
Tree tot.	ODW-g	$\ln Y = 1.678 + 2.352 \ln H_1$	10	.95	0.0 - 75 (mm)
<u>Rhododendron</u>					
(Telfer, 1969)					
Tree	ODW-g	$\ln Y = -2.833 + 2.612 \ln D_4$	20	.92	2.8 - 10.7 (mm)
Foliage	ODW-g	$\ln Y = -3.075 + 2.050 \ln D_4$	20	.83	2.8 - 10.7 (mm)
<u>Salal</u>					
(Stanek and State, 1978)					
Tree	ODW-g/m ²	$Y = 3.612 V + .02464 V^2$	50	.92	- - -
Tree	ODW-g/m ²	$Y = 28.65 ME$	35	.90	- - -
<u>Spirea</u>					
Meadow Sweet					
(Telfer, 1969)					
Tree	ODW-g	$\ln Y = -2.337 + 2.579 \ln D_4$	20	.91	1.0 - 12.6 (mm)
Foliage	ODW-g	$\ln Y = -2.257 + 1.720 \ln D_4$	20	.90	1.0 - 12.6 (mm)
<u>Spruce</u>					
Black					
(Keen, 1963)					
Bole, s*, t 3", wood	FW-lbs	$\log Y = 0.321 + 2.430 \log D_1$	58	.92 (r)	10.0 - 40.0 (cm)
Bole, s*, t 3"	FW-lbs	$\log Y = 0.477 + 2.345 \log D_1$	318	.95 (r)	10.0 - 40.0 (cm)
Tree, s*	FW-lbs	$\log Y = 0.755 + 2.209 \log D_1$	132	.96 (r)	10.0 - 40.0 (cm)
Bole, s*, t 3"	FW-lbs	$\log Y = -0.5474 + 1.861 \log D_1 + 0.839 \log H_1$	312	.98 (r)	10.0 - 40.0 (cm)
Bole, s*, t 3", wood	FW-lbs	$\log Y = -1.0122 + 1.724 \log D_1 + 1.142 \log H_1$	57	.98 (r)	10.0 - 40.0 (cm)
Slash	FW-lbs	$\log Y = 0.500 + 1.865 \log D_1$	132	.81 (r)	10.0 - 40.0 (cm)

Appendix

SPECIES (Reference)	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Spruce</u>					
Norway (Hakkila, 1972)					
Stump * and Roots 5 cm * height = 2.6 + 0.26 diameter of stump top	ODW-kg	$Y = -7.1 + 0.060 S_1^2$	239	.83 (r)	15 - 40 (cm)
<u>Spruce</u>					
Norway (Hakkila, 1972)					
Slash	FW-kg	$Y = -7.8 + 2.03 D_2 + 0.169 D_2$	245	.88 (r)	5 - 25 (cm)
Slash	ODW-kg	$Y = 5.9 + 0.00122 D_2^2 CR^2$	245	.93 (r)	10 - 30 (cm)
Foliage	ODW-kg	$Y = -16.0 + 5.18 D_2 - 0.0634 D_2 L + 0.118 CR$	245	.83 (r)	7.0 - 30.0 (cm)
Branches, wood	ODW-kg	$Y = 2.9 - 1.01 D_2 + 0.055 D_2^2 + 0.16 D_2 BR$	245	.94 (r)	10 - 30 (cm)
Slash, t < 6 cm, wood	ODW-kg	$Y = 4.4 - 0.99 D_2 + 0.0557 D_2^2 + 0.16 D_2 BR$	245	.93 (r)	10 - 30 (cm)
<u>Spruce</u>					
Red (Young, et al., 1964)					
Tree tot.	FW-lbs	$\ln Y = .6972 + 2.2234 \ln D_1 + .3390 \ln H_1$	25	.98	14.3 - 24.5 (cm)
Tree tot.	ODW-lbs	41% of FW from above equation	25	.98	14.3 - 24.5 (cm)
Tree tot. - Roots < 4"	FW-lbs	$\ln Y = 1.1400 + 2.2608 \ln D_1 + .1933 \ln H_1$	25	.98	14.3 - 24.5 (cm)
Tree tot. - Roots < 4"	ODW-lbs	41% of FW from above equation	25	.98	14.3 - 24.5 (cm)
Tree a.s. 6"	FW-lbs	$\ln Y = .1922 + 2.1679 \ln D_1 + .4292 \ln H_1$	25	.98	14.3 - 24.5 (cm)
Tree a.s. 6"	ODW-lbs	40% of FW from above equation	25	.98	14.3 - 24.5 (cm)
Stem a.s. 6"	FW-lbs	$\ln Y = -1.5151 + 1.8801 \ln D_1 + .9332 \ln H_1$	25	.99	14.3 - 24.5 (cm)
Stem a.s. 6"	ODW-lbs	49% of FW from above equation	25	.99	14.3 - 24.5 (cm)
Bole a.s. 6", t. 4"	FW-lbs	$\ln Y = -2.0267 + 2.0215 \ln D_1 + .9670 \ln H_1$	25	.99	14.3 - 24.5 (cm)
Bole a.s. 6", t. 4"	ODW-lbs	50% of FW from above equation	25	.99	14.3 - 24.5 (cm)
All branches	FW-lbs	$\ln Y = 3.1780 + 3.3292 \ln D_1 - 1.2874 \ln H_1$	25	.91	14.3 - 24.5 (cm)
All branches	ODW-lbs	10% of FW from above equation	25	.91	14.3 - 24.5 (cm)
Branches, < 1 in	FW-lbs	$\ln Y = 3.4430 + 2.9884 \ln D_1 - 1.2006 \ln H_1$	25	.88	14.3 - 24.5 (cm)
Branches, < 1 in	ODW-lbs	9% of FW from above equation	25	.88	14.3 - 24.5 (cm)

Appendix

SPECIES (Reference)	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Spruce</u>					
<u>Red</u>					
(Young, et al., 1964)					
.... cont'd					
Stump 6" and roots > 1 in	FW-lbs	$\ln Y = -.8508 + 2.5050 \ln D_1 + .1672 \ln H_1$	25	.95	14.3 - 24.5 (cm)
Stump 6" and roots > 1 in	ODW-lbs	45% of FW from above equation	25	.95	14.3 - 24.5 (cm)
Roots > 1 in	FW-lbs	$\ln Y = -.6402 + 2.6671 \ln D_1 + .0441 \ln H_1$	25	.93	14.3 - 24.5 (cm)
Roots > 1 in	ODW-lbs	44% of FW from above equation	25	.93	14.3 - 24.5 (cm)
Roots < 1 in	FW-lbs	$\ln Y = -1.5671 + 1.6306 \ln D_1 + .3564 \ln H_1$	25	.71	14.3 - 24.5 (cm)
Roots < 1 in	ODW-lbs	40% of FW from above equation	25	.71	14.3 - 24.5 (cm)
<u>Spruce</u>					
<u>Red sp.</u>					
(Young and Carpenter, 1967)					
Foliage	FW-g	$\ln Y = 3.674 + 1.619 \ln H_1$	40	.94	2.5 - 10.0 (cm)
Branches	FW-g	$\ln Y = 2.354 + 1.836 \ln H_1$	40	.95	2.5 - 10.0 (cm)
Stem	FW-g	$\ln Y = 1.242 + 2.687 \ln H_1$	40	.98	2.5 - 10.0 (cm)
Roots	FW-g	$\ln Y = 2.070 + 2.145 \ln H_1$	40	.97	2.5 - 10.0 (cm)
Tree tot.	FW-g	$\ln Y = 3.972 + 2.031 \ln H_1$	40	.98	2.5 - 10.0 (cm)
Foliage	ODW-g	$\ln Y = 2.891 + 1.626 \ln H_1$	40	.94	2.5 - 10.0 (cm)
Branches	ODW-g	$\ln Y = 1.833 + 1.847 \ln H_1$	40	.95	2.5 - 10.0 (cm)
Stem	ODW-g	$\ln Y = 1.512 + 2.705 \ln H_1$	40	.98	2.5 - 10.0 (cm)
Roots	ODW-g	$\ln Y = 1.376 + 2.136 \ln H_1$	40	.97	2.5 - 10.0 (cm)
Tree tot.	ODW-g	$\ln Y = 3.239 + 2.055 \ln H_1$	40	.98	2.5 - 10.0 (cm)
<u>Spruce</u>					
<u>White</u>					
(Baskerville, 1965)					
Tree	FW-lbs	$\log Y = 0.415 + 2.64 \log D_1$	12	.99	2.5 - 25.0 (cm)
Tree	ODW-lbs	$\log Y = 0.150 + 2.48 \log D_1$	13	.99	2.5 - 25.0 (cm)
Stem wood	ODW-lbs	$\log Y = 0.028 + 2.36 \log D_1$	13	.99	2.5 - 25.0 (cm)
Stem bark	ODW-lbs	$\log_{100} Y = 0.885 + 2.61 \log_{100} D_1$	14	.99	2.5 - 25.0 (cm)

Appendix

SPECIES (Reference) Equation Title	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Spruce</u> <u>White</u> (Baskerville, 1965) cont'd					
Branches	ODW-lbs	$\log Y = -0.855 + 2.78 \log D_1$	14	.97	2.5 - 25.0 (cm)
Foliage	ODW-lbs	$\log_{10} Y = 0.066 + 2.85 \log_{10} D_1$	14	.97	2.5 - 25.0 (cm)
Dead branches	ODW-lbs	$\log_{10} Y = -0.175 + 2.49 \log_{10} D_1$	13	.94	2.5 - 25.0 (cm)
<u>Spruce</u> <u>White</u> (Keen, 1963)					
Bole, s*, t 3", wood	FW-lbs	$\log Y = 0.090 + 2.601 \log D_1$	50	.99 (r)	10.0 - 52.5 (cm)
Bole, s*, t 3"	FW-lbs	$\log Y = 0.370 + 2.425 \log D_1$	164	.98 (r)	10.0 - 52.5 (cm)
Tree, s*	FW-lbs	$\log Y = 0.759 + 2.217 \log D_1$	112	.98 (r)	10.0 - 52.5 (cm)
Bole, s*, t 3"	FW-lbs	$\log Y = -0.4907 + 1.844 \log D_1 + 0.831 \log H_1$	160	.99 (r)	10.0 - 52.5 (cm)
Bole, s*, t 3", wood	FW-lbs	$\log Y = -0.5643 + 1.854 \log D_1 + 0.929 \log H_1$	47	.99 (r)	10.0 - 52.5 (cm)
Slash	FW-lbs	$\log Y = 0.742 + 1.753 \log D_1$	112	.82 (r)	10.0 - 52.5 (cm)
* = D ₁					
<u>Spruce</u> <u>White</u> (Kilil, 1967)					
Branches - live	ODW-lbs	$\log Y = -0.51780 + 1.27747 \log D_1 + 1.35248 \log CW$	(4)	.95	4.5 - 60.8 (cm)
Crown	ODW-lbs	$\log Y = -0.384118 + 1.35610 \log D_1 + 1.19533 \log CW$	60	.97	4.5 - 66.8 (cm)
<u>Spruce</u> <u>White</u> (Kimmins, 1974)					
Bole, a.s. 30.5 cm, t. 2.5 cm, wood	ODW-kg	$\log Y = 2.152 + 0.929 \log D_7$	11	.97	14.1 - 90.0 (cm)
Bole, a.s. 30.5 cm, t. 2.5 cm, bark	ODW-kg	$\log Y = 3.006 + 2.253 \log D_2$	11	.95	14.1 - 90.0 (cm)
Branches > 2.5 cm	ODW-kg	$\log Y = -4.863 + 3.256 \log (BA + CL_m CW_m)$	11	.76	14.1 - 90.0 (cm)
Branches, .64 - 2.5 cm	ODW-kg	$\log Y = 0.903 + 0.007 \log (BA + CL_m CW_m)$	11	.77	14.1 - 90.0 (cm)
Branches, < .64 cm	ODW-kg	$\log Y = 0.795 + 0.054 \log D_3 CL_m$	11	.87	14.1 - 90.0 (cm)
Foliage	ODW-kg	$\log Y = 1.044 + 0.053 \log D_3 CL_m$	11	.84	14.1 - 90.0 (cm)

Appendix

SPECIES (Reference) Equation Title	FW or ODW	PREDICTION EQUATION	n	R ² or r	RANGE OF DIAMETERS AT GROUND LEVEL IN MM OR DIAMETER AT BREAST HEIGHT IN CM
<u>Willow</u> (Ribe, 1973)					
Foliage	FW-g	$\log Y = 2.5653 + 1.6978 \log D_2$	30	.68	2.5 - 7.6 (cm)
Branches	FW-g	$\log Y = 2.6951 + 1.7736 \log D_2$	30	.70	2.5 - 7.6 (cm)
Stem	FW-g	$\log Y = 3.0186 + 2.4204 \log D_2$	30	.93	2.5 - 7.6 (cm)
Foliage	ODW-g	$\log Y = 2.1879 + 1.6442 \log D_2$	30	.67	2.5 - 7.6 (cm)
Branches	ODW-g	$\log Y = 2.4822 + 1.6624 \log D_2$	30	.77	2.5 - 7.6 (cm)
Stem	ODW-g	$\log Y = 2.7610 + 2.3391 \log D_2$	30	.91	2.5 - 7.6 (cm)
<u>Willow</u> (Telfer, 1969)					
Tree	ODW-g	$\ln Y = -1.519 + 2.325 \log D_2$	20	.97	4.5 - 34.1 (mm)
Foliage	ODW-g	$\ln Y = -1.324 + 1.729 \log D_4$	20	.82	4.5 - 34.1 (mm)
<u>Witch Hazel</u> (Telfer, 1969)					
Tree	ODW-g	$\ln Y = -3.037 + 2.900 \ln D_4$	21	.99	1.8 - 43.1 (mm)
Foliage	ODW-g	$\ln Y = -2.729 + 2.162 \ln D_4$	20	.96	1.8 - 43.1 (mm)
<u>White Red</u> (Telfer, 1969)					
Tree	ODW-g	$\ln Y = -2.613 + 2.774 \ln D_4$	20	.98	2.9 - 31.0 (mm)
Foliage	ODW-g	$\ln Y = -3.111 + 2.205 \ln D_4$	20	.95	2.9 - 31.0 (mm)